


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C07D 413/06, A61K 31/445 C07D 498/04, 401/06, 417/06 C07D 261/20, 413/12, 211/32</p>	A1	<p>(11) International Publication Number: WO 92/17475</p> <p>(43) International Publication Date: 15 October 1992 (15.10.92)</p>
<p>(21) International Application Number: PCT/US92/01605</p> <p>(22) International Filing Date: 9 March 1992 (09.03.92)</p> <p>(30) Priority data: 676,918 28 March 1991 (28.03.91) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 676,918 (CIP) Filed on 28 March 1991 (28.03.91)</p> <p>(71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).</p>		<p>(72) Inventors; and (75) Inventors/Applicants (for US only) : VILLALOBOS, Anabella [US/US]; 47 Greencliff Drive, Niantic, CT 06357 (US). NAGEL, Arthur, Adam [US/US]; 59 Inchcliffe Drive, Gales Ferry, CT 06335 (US). CHEN, Yuhpyng, Liang [US/US]; 8 Waterview Drive, Waterford, CT 06385 (US).</p> <p>(74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).</p> <p>(81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), CS, DE (Utility model), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), NO, PL, RU, SE (European patent), US.</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: HETEROCYCLIC-CYCLIC AMINE DERIVATIVES</p> <div style="text-align: center; margin: 20px 0;">  <p>(I)</p> </div> <p>(57) Abstract</p> <p>Compounds of formula (I) wherein R¹, R², R⁷, R⁸, X, Y, M and L are defined as below. The compounds of formula (I) are cholinesterase inhibitors and are useful in enhancing memory in patients suffering from dementia and Alzheimer's disease.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

-1-

5

HETEROCYCLIC-CYCLIC AMINE DERIVATIVESBackground of the Invention

The present invention relates to heterocyclic-cyclic amine derivatives of the formula I below, and pharmaceutically acceptable salts of such compounds. The
10 compounds of formula I are cholinesterase inhibitors and are useful in enhancing memory in patients suffering from dementia and Alzheimer's disease.


Alzheimer's disease is associated with degeneration of cholinergic neurons in the basal forebrain that play a
15 fundamental role in cognitive functions, including memory. Becker et al., Drug Development Research, 12, 163-195 (1988). As a result of such degeneration, patients suffering from the disease exhibit a marked reduction in acetylcholine synthesis, choline acetyltransferase activity,
20 acetylcholinesterase activity and choline uptake.

It is known that acetylcholinesterase inhibitors are effective in enhancing cholinergic activity and useful in improving the memory of Alzheimer's patients. By inhibiting acetylcholinesterase enzyme, these compounds increase the
25 level of the neurotransmitter acetylcholine, in the brain and thus enhance memory. Becker et al., supra, report that behavioral changes following cholinesterase inhibition appear to coincide with predicted peak levels of acetylcholine in the brain. They also discuss the efficacy
30 of the three known acetylcholinesterase inhibitors physostigmine, metrifonate, and tetrahydroaminoacridine.

-2-

Summary of the Invention

The present invention relates to compounds of the formula



10

wherein R^1 and R^2 are independently selected from hydrogen, (C_1-C_6) alkoxy, benzyloxy, phenoxy, hydroxy, phenyl, benzyl, halo, nitro, cyano, COR^5 , $-COOR^5$, $-CONHR^5$, $-NR^5R^6$, $-NR^5COR^6$, $-OCONR^5R^6$, $-NHCOOR^5$, (C_1-C_6) alkyl optionally substituted with
 15 from 1 to 3 fluorine atoms; SO_pCH_2 -phenyl or $SO_p(C_1-C_6)$ alkyl, wherein p is 0, 1 or 2; pyridylmethyloxy or thienylmethyloxy; 2-oxazolyl, 2-thiazolyl and benzenesulfonamide; wherein the phenyl moieties of said phenoxy, benzyloxy, phenyl, benzyl and benzenesulfonamide
 20 groups, the pyridyl and thienyl moieties of said pyridylmethyloxy or thienylmethyloxy and the oxazolyl and thiazolyl moieties of said 2-oxazolyl and 2-thiazolyl may optionally be substituted with 1 or 2 substituents independently selected from halo, (C_1-C_4) alkyl, trifluoromethyl, (C_1-C_4) alkoxy, cyano, nitro and hydroxy.
 25


or R^1 and R^2 , when attached to adjacent carbon atoms and when X is oxygen, sulfur or NR^4 wherein R^4 is hydrogen or (C_1-C_4) alkyl) may form, together with the carbon atoms to which they are attached, a group of the formula

30

35

-3-

5



10 wherein J is oxygen, sulfur or NR^4 , "a" is 1 or 2, R^3 is hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$ and Q is oxygen, sulfur, NH, CHCH_3 , $\text{C}(\text{CH}_3)_2$, $-\text{CH}=\text{CH}-$, or $(\text{CH}_2)_1$, wherein 1 is an integer from 1 to 3;

X is oxygen, sulfur, $-\text{CH}=\text{CH}-$, $-\text{CH}=\text{N}-$, $-\text{N}=\text{CH}-$, $-\text{N}=\text{N}-$, or
 15 NR^4 wherein R^4 is hydrogen or $(\text{C}_1\text{-C}_4)\text{ alkyl}$;


Y is $-(\text{CH}_2)_m-$, $-\text{CH}=\text{CH}(\text{CH}_2)_n-$, $-\text{NR}^4(\text{CH}_2)_m-$, or $-\text{O}(\text{CH}_2)_m-$
 wherein R^4 is defined as above, n is an integer from
 0 to 3 and m is an integer from 1 to 3;

R^5 and R^6 are each independently selected from hydrogen,
 20 $(\text{C}_1\text{-C}_6)\text{alkyl}$, phenyl or benzyl, wherein the phenyl moieties
 of said phenyl and benzyl may optionally be substituted with
 1 or 2 substituents independently selected from fluoro,
 chloro, bromo, iodo, $(\text{C}_1\text{-C}_4)\text{ alkyl}$, trifluoromethyl, $(\text{C}_1\text{-C}_4)$
 alkoxy, cyano, nitro and hydroxy, or NR^5R^6 together form a 4
 25 to 8 membered ring wherein one atom of the ring is nitrogen
 and the others are carbon, oxygen or nitrogen (e.g.
 pyrrolidinyl, piperidinyl, morpholino, piperazinyl or N-
 methylpiperazinyl), or NR^5COR^6 together form a 4 to 8
 membered cyclic lactam ring;

30 M is $-\text{CH}-$ or nitrogen;

L is phenyl, phenyl- $(\text{C}_1\text{-C}_6)\text{alkyl}$, cinnamyl or
 pyridylmethyl, wherein the phenyl moieties of said phenyl
 and phenyl- $(\text{C}_1\text{-C}_6)\text{alkyl}$ may optionally be substituted with 1
 to 3 substituents independently selected from $(\text{C}_1\text{-C}_6)\text{alkyl}$,
 35 $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_1\text{-C}_4)\text{alkoxycarbonyl}$, $(\text{C}_1\text{-C}_4)\text{alkylcarbonyl}$,
 $-\text{OCONR}^5\text{R}^6$, $-\text{NHCOOR}^5$ or halo; or L is a group of the formula

-4-



5

wherein b is an integer from 1 to 4, R^{13} and R^{14} are independently selected from hydrogen, (C_1-C_4) alkyl, halo and phenyl, E and F are independently selected from -CH- and
10 nitrogen, and G is oxygen, sulfur or NR^4 wherein R^4 is defined as above, with the proviso that when E and F are both nitrogen, one of R^{13} and R^{14} is absent; and

R⁷ and R⁸ are independently selected from hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkylcarbonyl and (C₁-C₆)alkoxy, with the proviso that said (C₁-C₆)alkoxy is not attached to a carbon that is adjacent to a nitrogen.

This invention also relates to the pharmaceutically acceptable acid addition salts of compounds of the formula I. Examples of such pharmaceutically acceptable acid addition salts are the salts of hydrochloric acid, p-toluenesulfonic acid, maleic acid, fumaric acid, citric acid, succinic acid, salicylic acid, oxalic acid, hydrobromic acid, phosphoric acid, methanesulfonic acid, tartaric acid, di-p-toluoyl tartaric acid, and mandelic acid.

This invention also relates to a pharmaceutical composition for inhibiting cholinesterase comprising a compound of the formula I or a pharmaceutically acceptable acid addition salt thereof, and a pharmaceutically acceptable carrier.


This invention also relates to a method for inhibiting cholinesterase in a mammal comprising administering to a mammal an amount of a compound of the formula I or a pharmaceutically acceptable acid addition salt thereof effective in inhibiting cholinesterase.

This invention also relates to a method for enhancing memory or treating or preventing Alzheimer's disease in a

-5-


mammal comprising administering to a mammal an amount of a compound of the formula I or a pharmaceutically acceptable acid addition salt thereof effective in enhancing memory or treating or preventing Alzheimer's disease.

5 This invention also relates to compounds of the formula



15 wherein W is a leaving group; j is an integer from 0 to 2; R¹⁰ is a nitrogen protecting group; and R⁷ and R⁸ are independently selected from hydrogen, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylcarbonyl and (C₁-C₆) alkoxy, with the proviso that said (C₁-C₆) alkoxy is not attached to a carbon that is adjacent to a carbon that is adjacent to a nitrogen. These compounds are useful as intermediates in the synthesis of compounds of the formula I.

This invention also relates to compounds of the formula




wherein R¹, R², R⁷, R⁸, X, Y and M are as defined above and R¹¹ is hydrogen or a nitrogen protecting group. These compounds are useful as intermediates in the synthesis of compounds of the formula I.

35

-6-

This invention also relates to compounds of the formula:



10

wherein R^1 , R^2 , R^7 , R^8 and L are as defined above and Y' is $-\text{CH}=\text{CH}-(\text{CH}_2)_n-$ or $-(\text{CH}_2)_m-$. These compounds are useful as intermediates in the synthesis of compounds of the formula I.

The term "mammal", as used herein, includes humans.

15 The term "halo", as used herein, includes chloro, bromo, iodo or fluoro.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched, or cyclic moieties or combinations thereof.

20 The term " (C_1-C_4) alkylcarbonyl", as used herein, refers to a substituent of the formula



25

wherein R^{15} is (C_1-C_4) alkyl.

30 The term " (C_1-C_4) alkoxy carbonyl", as used herein, refers to a substituent of the formula V above, wherein R^{15} is (C_1-C_4) alkoxy.

The term " (C_1-C_6) alkoxy carbonyl", as used herein, refers to a substituent of the formula V above, wherein R^{15} is (C_1-C_6) alkoxy.

35 The term " (C_1-C_6) alkylcarbonyl", as used herein, refers to a substituent of the formula V above, wherein R^{15} is (C_1-C_6) alkyl.

-7-

Preferred compounds of this invention are compounds of the formula I wherein X is oxygen or sulfur, Y is -CH₂-, -CH₂-CH₂-, M is -CH- and L is benzyl, R¹ and R² are (C₁-C₆)alkyl, (C₁-C₆)alkoxy, NR⁵R⁶, or NR⁵COR⁶, R³ is hydrogen or
5 (C₁-C₆)alkyl, J is oxygen or sulfur and Q is CH(CH₃), CH(CH₃)₂, -CH=CH or (CH₂)₁ and the pharmaceutically acceptable salts of such compounds.

3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

10 5-Methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5,6-Dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

15 5-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

7-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

20 6-Acetamido-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-Amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

25 6-Benzamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-Benzenesulfonamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-(4-Morpholinyl)-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

30 5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;

1-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-isoquinoline;

35 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisothiazole;

4-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,3-quinazoline;

-8-

- 6-Hydroxy-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;
- 6-Bromo-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;
- 5 6-Cyano-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;
- 6-Carboxamide-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;
- 3-[(1-Phenylmethyl-4-piperidyl)methoxy]-1,2-benzisoxazole;
- 10 3-[(1-Phenylmethyl-4-piperidyl)methylamino]-1,2-benzisoxazole;
- 3-[2-(1-Phenylmethyl)-4-piperidyl]ethylamino]-1,2-benzisoxazole;
- 15 3-[3-[1-(Phenylmethyl)-4-piperidyl]propyl]-1,2-benzisoxazole;
- trans-3-[2-[1-(Phenylmethyl)-4-piperidyl]ethenyl]-1,2-benzisoxazole;
- 3-[2-[1-(Phenylmethyl)-4-piperazinyl]ethyl]-1,2-benzisoxazole;
- 20 5,7-Dihydro-7-methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 5,7-Dihydro-7-ethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 25 5,7-Dihydro-3-[2-[1-(2-chloro-5-thiophenemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 5,7-Dihydro-3-[2-[1-(2-methyl-4-thiazolemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 3- [2-[1-(3-Bromophenylmethyl)-4-piperidinyl]-5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 30 3-[2-[1-(4-Bromophenylmethyl)-4-piperidinyl]ethyl]-5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 5,7-Dihydro-3-[3-[1-(phenylmethyl)-4-piperidinyl]propyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 35 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-5,6,8-trihydro-7H-isoxazolo[4,5-g]quinolin-7-one;

-9-

6,8-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-7H-pyrrolo[5,4-g]-1,2-benzisoxazol-7-one;

5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-6H-pyrrolo[5,4-f]-1,2-benzisoxazol-6-one;

5 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1H-indazole;

and the pharmaceutically acceptable salts of such compounds.

Examples of other compounds of the formula I are:

10 6-Phenylamino-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole;

6-(2-Thiazolyl)-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole;

6-(2-Oxazolyl)-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole;

15 6-Pyrrolidinyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole;

6-Piperidinyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole;

20 5,7-Dihydro-5,5-dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;

5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-7-n-propyl-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;

5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-7-i-propyl-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;

25 5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;

3[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-6-phenylmethanesulfone-1,2-benzisoxazole;

30 1-Methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1H-indazole; and

3-[1-Phenylmethyl-4-piperidinyl)methyl]-1,2-benzisoxazole.

The compounds of formula I may have optical centers and
35 may therefore occur in different isomeric forms. The invention includes all stereoisomers of such compounds of formula I, including mixtures thereof.


-10-

Detailed Description of the Invention


The preparation of compounds having the formula I and certain of the starting materials used in their synthesis is illustrated in the following reaction schemes. Except where
5 otherwise stated, in the reaction schemes and discussion that follow, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R¹³, R¹⁴, E, G, X, Y, M, L, a, b, l, m, n, p, and structures I, A, B, and K are defined as above.


All articles, books, patents and patent applications
10 cited in the following discussion are incorporated herein by reference.

-11-


SCHEME 1

-12-


SCHEME 2

SCHEME 2'

-14-


SCHEME 3

-15-


SCHEME 4

-16-


SCHEME 5



-17-

SCHEME 6

-18-



-19-

5 The preparation of compounds of the formula I wherein Y is $-(CH_2)_m$ and M is $-CH-$ is illustrated in scheme 1. These compounds are designated in scheme 1 and hereinafter referred to as compounds of the formula I-A (those wherein L is phenyl- (C_1-C_6) alkyl, pyridylmethyl or a group of the
10 formula K) and compounds of the formula I-B (those wherein L is phenyl or cinnamyl).

Referring to scheme 1, compounds of the formula I-A may be prepared by deprotonating a compound of the formula II with a base in the presence of, or followed by the addition
15 of an alkylating agent of the formula III wherein R^{10} is a nitrogen protecting group and W is a leaving group. When R^{10} is a nitrogen protecting group, this reaction produces an intermediate of the formula IV. This intermediate is then deprotected to yield a secondary piperidine of the formula
20 VI as a free base or a salt of the free base, after which such free base or salt is alkylated with a compound of the formula WL, wherein W is defined as above and L is phenyl- (C_1-C_6) alkyl, pyridylmethyl or a group of the formula K.

Examples of suitable leaving groups (W) are mesylate,
25 tosylate, chloride, iodide and bromide. Examples of suitable nitrogen protecting groups (R^{10}) are amides such as N-formyl and N-acetyl and carbamates such as t-butoxycarbamate (BOC). The preferred nitrogen protecting group is BOC. Appropriate bases for use in the preparation
30 of compounds of the formula IV include strong bases such as lithium diisopropylamide (LDA), n-butyllithium, s-butyllithium, and lithium (or sodium or potassium) hexamethyldisilazide (LiHMDS, NaHMDS, or KHMDS). LDA and s-butyllithium are preferred.

35 The reaction of a compound of formula II with a compound of formula III is generally carried out in a polar, aprotic solvent such as diethyl ether, 1,2-dimethoxyethane, or tetrahydrofuran (THF). Temperatures may range from about -78°C to about 30°C . This reaction is preferably conducted
40 in THF at about -78°C .

-20-

Usually, compounds of the formula II are deprotonated in the presence of a compound of formula III. However, in cases where the compound of formula II has more than one acidic proton, it is preferable to carry out the deprotonation step first followed by the immediate and rapid addition of the alkylating agent of formula III.

The protecting group (R^{10}) can be removed from compounds of the formula IV to form the corresponding compounds of formula VI by methods known to those skilled in the art.

10 For example, when R^{10} is BOC or another carbamate, it can be removed with an acid such as hydrogen bromide (gas or aqueous), hydrogen chloride (gas or aqueous) or trifluoroacetic acid. In the case of trifluoroacetic acid, a t-butyl cation scavenger such as thioanisole may be added.

15 When an acid is used as the deprotecting agent, an acid addition salt of the compound of formula VI is produced rather than the free base of such compound. Appropriate solvents include non-polar solvents such as methylene chloride, as well as polar solvents such as diethyl ether,

20 ethyl acetate, dioxane, alcohols (e.g. methanol or ethanol) and water. Temperatures may range from about -20°C to about the reflux temperature of the solvent. It is preferable to use trifluoroacetic acid in methylene chloride with or without thioanisole at about 0°C .

25 Alternatively, when R^{10} is BOC, it can be removed with a trialkylsilyltrifluoromethanesulfonate derivative such as trimethylsilyl-, triethylsilyl-, or t-butyldimethylsilyltrifluoromethanesulfonate in the presence of an aromatic or tertiary amine base such as 2,6-lutidine or triethylamine.

30 Appropriate solvents for this reaction include nonpolar solvents such as methylene chloride and polar aprotic solvents such as THF, diethyl ether or DMF. Temperatures may range from about -20°C to room temperature. It is preferable to use trimethylsilyltrifluoromethane-sulfonate

35 and 2,6-lutidine in methylene chloride at a temperature from about 0°C to about room temperature.

-21-

The intermediate secondary piperidine of the formula VI, obtained as the free base or salt as described above, is reacted with 2-10 equivalents of a base and then with an alkylating agent of the formula WL, wherein W is defined as
5 above and L is phenyl-(C₁-C₆)alkyl, pyridylmethyl or a group of the formula K. Suitable bases include tertiary amines such as triethylamine and diisopropylethyl-amine, aromatic amines such as pyridine and dimethylaminopyridine, and metal carbonates such as sodium bicarbonate or sodium or potassium
10 or cesium carbonate. When W is chloride, catalytic iodide (potassium iodide or tetra-n-butylammonium iodide) may be added. Appropriate solvents include non-polar solvents such as methylene chloride and polar solvents such as dimethylformamide, THF, acetonitrile, acetone, dioxane, and
15 alcohols such as methanol or ethanol. It is preferable to carry out the alkylation in the presence of triethylamine in methylene chloride at room temperature or in the presence of sodium carbonate in dimethylformamide at room temperature.

Alternatively, the intermediate secondary piperidine of
20 formula VI, when obtained as a salt after removal of the protecting group, can be deprotonated to the free amine by dissolving or suspending it in an appropriate solvent (e.g., methylene chloride or ethyl acetate), mixing it with aqueous sodium bicarbonate or aqueous sodium or potassium hydroxide
25 and recovering the free amine from the organic layer by conventional extraction techniques. The free amine can then be subjected to alkylation with the appropriate alkylating agent of the formula WL under the conditions described above using 1-2 equivalents of an appropriate base.

30 The starting materials of formula II can be prepared according to methods known in the art. When X is oxygen, the starting 3-methyl-1,2-benzisoxazoles can be prepared by procedures similar to those described by Wunsch *et al.*, Adv. Heterocycl. Chem., 1967, 8, 277; Smalley, R.K., Adv. Heterocycl. Chem. 1981, 29, 2; and Thakar *et al.*, Indian J. Chem. 1977, 15B, 1058. The appropriate o-hydroxy acetophenones are converted to the corresponding oximes by
35

-22-

reaction with hydroxylamine hydrochloride in the presence of an appropriate base such as potassium or sodium hydroxide, sodium acetate or pyridine, preferably aqueous potassium hydroxide or aqueous sodium acetate, in a polar solvent such as methanol, ethanol, or water, preferably ethanol, at a temperature from about room temperature to about 120°C. The oxime is then converted to the corresponding oxime acetate by acetylation with an appropriate acylating agent such as acetic anhydride. Temperatures for this reaction may range from about room temperature to the reflux temperature of the solvent. Temperatures between 80°C and 130°C are preferred.

Ring closure to form the benzisoxazole ring may be carried out by heating the neat oxime acetate at a temperature from about 125°C to about 200°C under atmospheric pressure or reduced pressure (e.g., from about 0.01 mm Hg to about 760 mm Hg). Ring closure is preferably accomplished by heating the oxime acetate at reflux in an appropriate base such as pyridine, or by heating the oxime acetate at a temperature of about 130°C in a polar solvent such as DMF or DMSO (dimethylsulfoxide) in the presence of several equivalents of an appropriate base such as pyridine or 2,6-lutidine.

Alternatively, ring closure can be carried out directly from the oxime by reaction with an acyl or sulfonyl chloride such as oxalyl or thionyl chloride in the presence of an aromatic amine such as pyridine (See Kalkote *et al.*, Aust. J. Chem. 1977, 30, 1847). Suitable solvents include polar solvents such as diethyl ether or THF. Temperatures can range from about 0°C to about room temperature. Another method of closure involves treatment of the oxime with one or less equivalents of a base such as potassium hydroxide in a polar solvent such as methanol at temperatures ranging from about room temperature to about 100°C (Crabbe *et al.*, J. Chem. Soc. Perkin Trans. I, 1973, 2220).

When X is sulfur, the starting 3-methyl-1,2-benzisothiazoles can be prepared from o-methylthio acetophenones following procedures similar to those

-23-

described above for the benzisoxazoles (See McKinnon et al., Can. J. Chem., 1988, 66, 1405 and references cited therein). The o-methylthio acetophenones are converted into the corresponding oximes and ring closure is carried out directly by reaction with an appropriate acylating agent such as acetic anhydride in a base such as pyridine. The reaction temperature may range from about room temperature to about 130°C, and is preferably about 120°C.

When X is NR⁴ wherein R⁴ is hydrogen the starting 3-methyl-1H-indazoles can be prepared according to methods described by Behr et al., "Pyrazoles, Pyrazolines, Pyrazolidines, Indazoles, and Condensed Rings," Heterocyclic Compounds, R.H. Wiley, Ed., 1967, 289; Bartsch et al., J. Heterocycl. Chem., 1984, 21, 1063; Hannig et al., Pharmazie 1976, 31, 534; Barton et al., J. Chem. Soc. Chem. Comm., 1982, 450; Ruechardt et al., Liebigs Ann. Chem., 1980, 908; and Rees et al., J. Chem. Soc. D, 1971, 827. N-Alkylation of 3-methyl-1H-indazoles (X is NR⁴ wherein R⁴ is (C₁-C₄)alkyl) can be carried out as described by Behr et al., "Pyrazoles, Pyrazolines, Pyrazolidines, Indazoles, and Condensed Rings," Heterocyclic Compounds, R. H. Wiley, Ed., 1967, 309; Palmer et al., J. Chem. Soc., Perkin Trans. II, 1975, 1695; and Claramunt et al., Heterocycles 1985, 23, 2895.

When X is -CH=CH-, the starting 1-methylisoquinolines can be prepared according to the Bischler-Napieralksi or Pictet-Spengler methods (See Organic Reactions, Vol. VI, chapters 2 and 3, pp. 74-190, John Wiley and Sons, New York, 1951).

When X is -N=CH-, the starting 4-methylquinazolines can be prepared according to methods described by Byford et al., Indian J. Chem., 1988, 27B, 396; Higashino, T., Chem. Pharm. Bull., 1962, 10, 1043; and Uff et al., J. Chem. Soc., Perkin Trans. I, 1986, 2295.

When X is -CH=N- the starting 1-methylphthalazines can be prepared according to methods described by Kant et al., J. Heterocycl. Chem., 1985, 22, 1065 and references cited


-24-

therein; Acheson et al., J.Chem. Soc. C, 1966, 2218; and Gabriel et al., Chem. Ber. 1897, 30, 3022.

When X is -N=N-, the starting 4-methyl-1,2,3-benzotriazines can be prepared according to methods
5 described by Adger et al., J. Chem. Soc., Perkin Trans. I, 1975, 31; Boulton et al., Ibid., 1988, 1509; and Rees et al., J. Chem. Soc. D, 1971, 828.

When one or both of R¹ and R² are NH₂, the starting material of formula II may be prepared from the
10 corresponding NHAc precursor (Ac=acetyl) by acid hydrolysis. The acid hydrolysis can be carried out with aqueous hydrochloric acid at temperatures ranging from about 50°C to about 120°C. Heating to reflux (about 120°C) in 1N HCl is preferred. The corresponding NHBz (Bz=benzoyl) or NHSO₂C₆H₅,
15 compounds can be prepared from the corresponding amino derivative by reaction with the appropriate benzoyl or benzenesulfonyl chloride in the presence of a base such as triethylamine, pyridine, or dimethylaminopyridine. Suitable solvents include methylene chloride, THF, diethylether, or
20 dimethylformamide. Temperatures may range from about -20° to about 80°C. When one or both of R¹ and R² is NHBz, it is preferable to use triethylamine/dimethylaminopyridine in methylene chloride at room temperature. When one or both of R¹ and R² is NHSO₂Ph, it is preferable to use pyridine in
25 methylene chloride at 0°C.

Cyclic dialkylamino compounds of the formula II (i.e., those wherein one or both of R¹ and R² are NR⁵R⁶ wherein NR⁵R⁶ together form a ring) can also be prepared from the
corresponding amino derivative by alkylation with the
30 appropriate bis-halide reagent of the formula




wherein each A is independently bromide or chloride and B is
35 oxygen or (CH₂)_q wherein q is 0 to 2, in the presence of an appropriate base such as triethylamine or diisopropylethylamine (Hunig's base), in an appropriate

-25-

nonpolar solvent such as toluene or xylene. The alkylation is typically carried out at a temperature from about room temperature to about 150°C. It is preferably conducted in the presence of Hunig's base in toluene at about 120°C (reflux). (See Verboom et al., J. Org. Chem. 1984, 49, 269).

Alternatively, these cyclic dialkylamino derivatives may be prepared by nucleophilic displacement of an aromatic fluoride with the appropriate cyclic amine. Suitable solvents for this reaction include polar aprotic solvents such as dimethylsulfoxide (DMSO), dimethylformamide (DMF), acetonitrile, pyridine and hexamethylphosphoramide. Acetonitrile and pyridine are preferred. The reaction may be run in the presence of a base such as a tertiary or aromatic amines (e.g., triethylamine, diisopropylethylamine, pyridine or dimethylaminopyridine), preferably pyridine or triethylamine. The reaction temperature may range from about room temperature to about 160°C, and is preferably from about 80°C to about 160°C.

20 When R^1 and R^2 , together with the carbons to which they
are attached, form a group of the formula




30

and X is oxygen or sulfur, the starting material of formula II may be prepared by the following procedure which is illustrated only for cases wherein R¹ and R² form a group of formula A.

35

-26-



First, a compound of the formula VIII wherein X' is hydroxy, thiol or methyl sulfide is prepared by Friedel-Crafts acylation of the corresponding compound of the formula VII, wherein X'' is methoxy or methyl sulfide with an acylating agent such as acetyl chloride or acetic anhydride, preferably acetyl chloride, in the presence of a Lewis acid such as aluminum chloride, titanium tetrachloride or boron trifluoride etherate, preferably aluminum chloride. Appropriate solvents include carbon disulfide, 1,2-dichloroethane and nitrobenzene. Carbon disulfide and 1,2-dichloroethane are preferred. Generally, this reaction is conducted at a temperature from about room temperature to about 200°C, preferably from about 50°C to about 100°C.

Alternatively, a compound of formula VIII wherein X' is hydroxy, may also be obtained by Fries rearrangement of the corresponding compound of the formula VII wherein X'' is acetyloxy. A mixture of VII and a Lewis acid such as aluminum chloride, boron trifluoride etherate or titanium tetrachloride is heated at temperatures from about 80°C to 200°C either neat or in the presence of a solvent such as nitrobenzene, or 1,2-dichloroethane. The Fries rearrangement is preferably conducted neat with aluminum chloride at 170-190°C.

The compounds of formula VIII obtained by the foregoing process may be converted to the corresponding starting materials of formula II by the procedure described above for 1,2-benzisoxazoles and 1,2-benzisothiazoles.

-27-

Referring to scheme 1, compounds of the formula III may be prepared from the corresponding compounds wherein W is hydroxy by methods known in the art. For example, compounds of the formula III wherein W is iodide can be prepared by
5 reacting the hydroxy counterpart with iodine and triphenylphosphine in the presence of a base such as pyridine or imidazole in a non-polar solvent such as benzene or toluene at a temperature from about room temperature to about 130°C. Preferably, the reaction is carried out in
10 benzene in the presence of pyridine at about 90°C (reflux).

Compounds of the formula I-B may be prepared by deprotonating a compound of the formula II with a base in the presence of, or followed by the addition of, an alkylating agent of the formula III wherein R¹⁰ is phenyl or
15 cinnamyl and W is defined as above. Suitable and preferred bases, solvents and conditions are similar to those described above for the preparation of compounds of the formula IV.

Scheme 2 illustrates the preparation of compounds of the formula I wherein Y is $-\text{CH}=\text{CH}(\text{CH}_2)_n-$ via an aldol-type
20 condensation. These compounds are designated in scheme 2 and hereinafter referred to as compounds of the formula I-C. Referring to scheme 2, a compound of the formula IX wherein R⁹ is hydrogen is deprotonated with a base followed by
25 immediate and rapid addition of an aldehyde of the formula X. Suitable bases and solvents are the same as those described above for the first reaction in scheme 1. The reaction temperature may range from about -78°C to about room temperature. The reaction is preferably carried out
30 using lithium diisopropylamide in THF at about -78°C and allowed to warm to room temperature.

If an intermediate alcohol is formed, it can be dehydrated to the olefin under standard acidic conditions, using an acid such as dilute hydrochloric acid, p-
35 toluenesulfonic acid or pyridinium p-toluenesulfonate, preferably p-toluenesulfonic acid, in a solvent such as benzene, toluene, THF or methylene chloride, at a

-28-

temperature from about 0°C to about 130°C. Preferably, the dehydration is carried out in benzene at about 80°C (reflux) with azeotropic removal of water. Dehydration may also be accomplished by treatment with Burgess' reagent

5 + -
(Et₃NSO₂NCO₂Me) in methylene chloride or benzene at a temperature from about room temperature to about 80°C.

Alternatively, the intermediate alcohol may be converted into a good leaving group such as mesylate or
10 tosylate and then eliminated with an appropriate base. The mesylate or tosylate can be prepared under standard conditions by reacting the alcohol with methanesulfonyl chloride or p-toluenesulfonyl chloride in the presence of a base such as triethylamine, diisopropylethylamine, or
15 pyridine. Appropriate solvents include methylene chloride and THF, with methylene chloride being preferred. Temperatures may range from about 0°C to about 60°C, and are preferably from about 0°C to about room temperature. Elimination to form the olefin can then be carried out with
20 a base such as diazabicycloundecane or diazabicyclononane in a suitable solvent such as benzene, methylene chloride, or THF, with benzene or methylene chloride being preferred, at a temperature from about 0°C to about 100°C, preferably from about room temperature to about 100°C.

25 Compounds of the formula I-C may also be prepared by a Wittig reaction from compounds of the formula IX wherein R⁹ is bromine, chlorine or iodine. According to this procedure, a compound of formula IX is converted into its phosphonium salt by treatment with triphenylphosphine in a
30 nonpolar solvent such as benzene, toluene or xylene, preferably toluene, at a temperature from about room temperature to about 150°, preferably from about 80°C to about 120°C. The phosphonium salt may then be deprotonated with a strong base such as sodium hydride, potassium t-
35 butoxide, potassium hydride or n-butyllithium in a suitable solvent such as diethylether or THF, at a temperature from about 0°C to about 80°C. The deprotonation is preferably

-29-

carried out with sodium hydride in THF at about room temperature.

Scheme 2' illustrates an alternative preparation of compounds of the formula I wherein X is oxygen or NR^4 , Y is $(\text{CH}_2)_m$ or $-\text{CH}=\text{CH}(\text{CH}_2)_n$ and M is carbon (i.e. $-\text{CH}-$). These compounds are designated in Scheme 2', and hereinafter referred to as compounds of the formula I-F (those where Y is $-\text{CH}=\text{CH}(\text{CH}_2)_n$ and n' is an integer from 0 to 3) and I-G (those where Y is $(\text{CH}_2)_m$ and n' is an integer from 0 to 1). Referring to Scheme 2', compounds of the formula I-F may be prepared by deprotonating a compound of the formula XV with a suitable base followed by addition of an aldehyde of formula XVI to give an intermediate of formula XVII. This intermediate is then transformed to compounds of the formula I-F by reaction with an appropriate amine.

Appropriate bases for use in the preparation of compounds of the formula XVII include lithium diisopropylamide, lithium or sodium or potassium hexamethyldisilazide, or n-butyllithium, preferably lithium diisopropylamide or lithium hexamethyldisilazide. The reaction of a compound of formula XV with a compound of formula XVI is generally carried out in a polar aprotic solvent such as diethyl ether, 1,2-dimethoxyethane, or tetrahydrofuran. Temperatures may range from -78°C to 80°C . This reaction is preferably conducted in THF at -78°C and allowed to warm to room temperature.

A compound of formula I-F is then obtained from an intermediate of formula XVII by reaction with an amine such as hydrazine or hydroxylamine in the presence of a base such as sodium or potassium hydroxide, sodium or potassium carbonate, or sodium or potassium alkoxide (methoxide or ethoxide), preferably sodium or potassium hydroxide. In some cases (when amine is hydrazine), addition of a base may not be necessary. Suitable solvents for this reaction include methanol, ethanol, i-propanol, water, or, when amine is hydrazine, hydrazine itself may be used as a solvent. Temperatures may range from 50°C to 120°C . It is preferable

-30-

to react XVII with hydrazine at 120°C (reflux), or with hydroxylamine and potassium hydroxide in EtOH/water at 100°C (reflux). Upon reaction with hydroxylamine, the intermediate oxime obtained may be isolated and then
5 cyclized to compounds of the formula I-F following the suitable and preferred conditions described above for the preparation of starting materials 3-methyl-1,2-benzioxazoles.

Compounds of the formula I-G may be prepared by
10 reducing an intermediate of formula XVII to give a compound of formula XVIII. The intermediate of formula XVII may then be reacted with an amine to give compounds of the formula I-G. An intermediate of formula XVII is reduced with hydrogen gas to an intermediate of formula XVIII in the presence of
15 a catalyst such as palladium on carbon, platinum oxide, or rhodium on carbon, preferably platinum oxide in a polar solvents such as ethyl acetate, tetrahydrofuran, ethanol, or acetic acid; preferably EtOH. Pressure may range from atmospheric to 50 psi, preferably 40-50 psi and temperature
20 may range from room temperature to 80°C, preferably room temperature.

A compound of formula I-G is then obtained from an intermediate of formula XVIII following the suitable and preferred conditions described above for the preparation a
25 compound of formula I-F from an intermediate of formula XVII.

Scheme 3 illustrates the preparation of compounds of the formula I wherein Y is $-O(CH_2)_m-$ or $-NR^4(CH_2)_m$. These compounds are designated in scheme 3 and hereinafter
30 referred to as compounds of the formula I-D. Referring to scheme 3, compounds of the formula I-D may be prepared by reacting a compound of the formula XI wherein R^{12} is chloro or bromo, with a nucleophile of the formula XII, wherein Z is $-NHR^4$ or $-OH$. When Y is $NR^4(CH_2)_m$ (i.e., Z is $-NHR^4$), the
35 amine of formula XII is generally reacted with the appropriate compound of formula XI either neat or in a polar solvent such as dimethylformamide (DMF), dimethylsulfoxide

-31-

(DMSO), THF or pyridine. DMSO and DMF are preferred solvents. An acid acceptor such as diazabicycloundecane, pyridine, lutidine, triethylamine or metal carbonates such as potassium or sodium or cesium carbonate may be added.

- 5 Metal carbonates such as potassium carbonate are preferred. The reaction temperature may range from about room temperature to about 160°C and is preferably from about 100°C to about 160°C.

When Y is $O(CH_2)_m$ (i.e., Z is -OH), the alkoxide anion
10 is formed and this species is reacted with the compound of formula XI. According to this procedure, the alcohol (XII) is deprotonated with a suitable base, after which the appropriate compound of formula XI is added and the mixture is heated. Examples of suitable bases are sodium, sodium
15 hydride and potassium hydride with sodium hydride being preferred. Suitable solvents include THF, DMF and DMSO, with THF and DMF being preferred. This reaction is generally conducted at a temperature from about 40°C to about 160°C. Temperatures between about 60°C and about
20 160°C are preferred.

Alternatively, compounds of the formula I-D may be prepared by reacting a compound of formula XI with a nucleophile of the formula XIX wherein Z and R^{10} are defined as above. This reaction produces an intermediate of formula
25 XX which is then deprotected to yield a secondary piperidine of formula XXI as a free base or a salt of the free base, after which such free base or salt is alkylated with a compound of the formula WL, wherein W is defined as above and L is phenyl(C_1 - C_6)alkyl, pyridylmethyl, or a group of the
30 formula K.

Suitable and preferred bases, solvents, and conditions for the reaction of a compound of formula XI with a nucleophile of the formula XIX are similar to those described for the reaction of a compound of the formula XI
35 with a nucleophile of the formula XII for the preparation of compounds of the formula I-D. Suitable and preferred bases, solvents, and conditions for the transformations of

-32-

compounds of the formulas XX and XXI to prepare compounds of the formula I-D are similar to those described in scheme 1 for compounds of the formulas IV and VI to prepare compounds of the formula I-A.

5 Scheme 4 illustrates the preparation of compounds of the formula I wherein Y is $-(CH_2)_m-$ and M is nitrogen. These compounds are designated in scheme 4 and hereinafter referred to as compounds of the formula I-E. Referring to scheme 4, a compound of the formula XIII wherein R^{12} is
10 chloro, bromo or iodo is reacted with a compound of the formula XIV. This reaction may be carried out in the presence of an acid acceptor such as pyridine, 2,6-lutidine or a metal carbonate (e.g., sodium bicarbonate or sodium or potassium carbonate). When R^{12} is chloro or bromo, a
15 catalytic amount of a displacement promoter may be added. Examples of suitable displacement promoters are sodium iodide, potassium iodide or tetra-n-butylammonium iodide. Generally, this reaction is conducted in a nonpolar solvent such as toluene or xylene, or in a polar solvent such as
20 THF, DMF or DMSO, preferably xylene or DMF, at a temperature from about room temperature to about 160°C, preferably from about 90°C to about 160°C.

Scheme 5 illustrates the preparation of compounds of the formula I wherein X is $-N=CH-$ and M is carbon (i.e.
25 $-CH$). These compounds are designated in scheme 5 and hereinafter referred to as compounds of the formula I-H. Referring to scheme 5, a compound of the formula II' may be deprotonated with one equivalent of a base followed by addition of a silylating agent (trimethylsilyl chloride).
30 Sequential deprotonation with a second equivalent of the same base followed by addition of an alkylating agent of the formula III and appropriate work-up produces an intermediate of the formula IV'. This intermediate is then deprotected as described in scheme I to yield a secondary piperidine of
35 formula VI' as a free base or a salt of the free base, after which such free base or salt is alkylated with a compound of the formula WL, wherein W is defined as above and L is

-33-

phenyl(C₁-C₆)alkyl, pyridylmethyl, or a group of the formula K.

Suitable bases, solvents, and temperatures for deprotonation of a compound of the formula II' are the same as those described above for the first reaction in scheme I, preferably LDA in THF at 0° to room temperature. After addition of the first equivalent of base, a silylating agent such as trimethylsilyl or triethylsilyl chloride is added, preferably trimethylsilyl chloride. A second equivalent of the same base is then added followed by an alkylating agent of the formula III. The trimethylsilyl group is then removed under acidic conditions by stirring the crude reaction mixture with dilute hydrochloric acid for 30-60 min at room temperature. Then, the crude reaction mixture is made basic with aqueous sodium carbonate, or aqueous sodium or potassium hydroxide, preferably aqueous sodium hydroxide, and the intermediate of formula IV' is extracted with an organic solvent by conventional extraction techniques. Suitable and preferred conditions for the transformation of intermediates of the formula IV' to compounds of the formula I-H are the same described in scheme I for the preparation of compounds of the formula I-A.

Scheme 6 illustrates the preparation of compounds of the formula I wherein Y is (CH₂)_m. M is carbon (i.e. -CH-), J is sulfur and Q is CHCH₃, C(CH₃)₂, -CH=CH, or (CH₂)₁. These compounds are designated in scheme 6 and hereinafter referred to as compounds of the formula I-I. Referring to scheme 6, a compound of the formula I-I may be prepared from the corresponding compound of the formula IV" where J is oxygen by reacting with a phosphorus sulfide to give an intermediate of the formula XXII. This intermediate is then deprotected as described in scheme I to yield a secondary piperidine of formula VI" as a free base or a salt of the free base, after which such free base or salt is alkylated with a compound of the formula WL, wherein W is defined as above and L is phenyl(C₁-C₆)alkyl, pyridylmethy, or a group of the formula K.

-34-

The transformation of a compound of formula IV" to an intermediate of formula XXII is carried out with a phosphorous sulfide such as phosphorous pentasulfide (P_2S_{10}) or Lawesson's reagent [2,4-bis(4-methoxyphenyl)-1,3-dithia-
5 2,4-diphosphetane-2,4-disulfide] in a non-polar solvent such as benzene, toluene, or xylene. Temperatures may range from 50°C to 160°C. Lawesson's reagent in toluene at 80°C is preferred. Suitable and preferred conditions for the transformation of intermediates of the formula XXII to
10 compounds of the formula I-I are the same ones described in scheme 1 for the preparation of compounds of the formula I-A.

Scheme 7 illustrates the preparation of compounds of the formula I wherein Y is $(CH_2)_m$, M is carbon (i.e. -CH-),
15 and R^3 is (C_1-C_6) alkyl. These compounds are designated in scheme 7 and hereinafter referred to as compounds of the formula I-J. Referring to scheme 7, a compound of the formula I-J may be prepared from the corresponding compounds of the formula IV''' where R^3 is hydrogen by deprotonating
20 with a base followed by addition of the appropriate alkylating agent (preferably the appropriate (C_1-C_6) alkyl chloride, bromide, or iodide) to give an intermediate of formula XXIII. This intermediate is then deprotected as described in scheme 1 to yield a secondary piperidine of
25 formula VI''' as a free base or a salt of the free base, after which such base or salt is alkylated with a compound of the formula WL, wherein W is defined as above and L is phenyl (C_1-C_6) alkyl, pyridylmethyl, or a group of the formula K.

30 Suitable bases for the transformation of a compound of the formula IV''' to a compound of the formula XXIII include sodium hydride, potassium hydride, lithium diisopropylamide, or n-butyllithium, preferably sodium hydride. The reaction is generally carried out in a polar aprotic solvent such as
35 tetrahydrofuran, dimethylformamide, or 1,2-dimethoxyethane and temperatures may range from -78°C to 80°C. The reaction is preferably conducted in dimethylformamide at room

-35-

temperature. Suitable and preferred conditions for the transformation of intermediates of the formula XXIII to compounds of the formula I-J are the same described in scheme 1 for the preparation of compounds of the formula I-A.

Alternatively, a compound of the formula I-J may be prepared directly from the corresponding compound of the formula I-A' wherein L is phenyl-(C₁-C₆) alkyl, pyridylmethyl or a group of the formula K and R³ is hydrogen or I-B' wherein L is phenyl or cinnamyl and R³ is hydrogen. Compounds of the formula I-A' and I-B' are prepared according to the methods described in scheme 1 for the preparation of compounds I-A and I-B. Suitable and preferred bases, solvents, and conditions for the transformation of compounds of the formula I-A' and I-B' to compounds of the formula I-J are the same described above for the preparation of compounds of the formula XXIII.

When one or both of R¹ and R² are OH, compounds of the formula I may be prepared from the corresponding -OMe precursor by dealkylation with a Lewis acid such as aluminum trichloride, boron trichloride, boron tribromide, or a protic acid such as aqueous hydrochloric or hydrobromic acid. Suitable solvents for the reaction with Lewis acids include non-polar solvents such as benzene, toluene, dichloromethane, or 1, 2-dichlorethane. Temperatures may range from -78°C to 120°C. Aqueous hydrobromic acid (48%) at 100-120°C (reflux) is preferred.

When one or both of R¹ and R² are NH₂, compounds of the formula I may be prepared from the corresponding NHAc precursor (Ac=acetyl) by acid hydrolysis under the suitable and preferred conditions described above for the formation of starting materials of formula II wherein one or both of R¹ and R² are NH₂. The corresponding nitrile (-CN) compounds can be prepared from the corresponding amino compounds via a diazonium salt formation by reacting the amino compound with nitrous acid (made from aqueous hydrochloric acid and sodium nitrite) followed by neutralization and addition to

-36-

CuCN. Suitable solvents include polar protic solvents such as water or biphasic mixtures with non-polar solvents such as benzene, toluene, or xylene. Neutralization may be carried out by adding a base such as sodium carbonate, 5 potassium carbonate, sodium hydroxide, or potassium hydroxide until pH 7. Temperatures may range from -20°C to 60°C. It is preferable to carry out the diazonium salt formation in water at 0°C, to neutralize with sodium carbonate, and to add the diazonium salt to a biphasic 10 mixture of aqueous CuCN and toluene at 0°C, followed by heating to 50°C.

When one or both of R¹ and R² are carboxamide (-CONH₂), compounds of the formula I may be prepared from the corresponding nitrile (-CN) precursor by reaction with a 15 base such as sodium hydroxide, potassium hydroxide or tetramethylammonium hydroxide in a polar solvent such as water, methanol, ethanol, or t-butanol. Temperatures may range from room temperature to 120°C. Potassium hydroxide in t-butanol at 85-100°C is preferred.

20 Compounds of the formula I other than those of formulae I-A-I-J may be prepared by methods that will be obvious to those skilled in the art from the procedures described above and other known methods.

In each of the above reactions, pressure is not 25 critical. Pressures in the range of about 0.5 atm to 3 atm are suitable, and ambient pressure (generally, about one atmosphere) is preferred as a matter of convenience. Also, for those reactions where the preferred temperature varies with the particular compounds reacted, no preferred 30 temperature is stated. For such reactions, preferred temperatures for particular reactants may be determined by monitoring the reaction using thin layer chromatography.

The compounds of formula I and their pharmaceutically acceptable salts (hereinafter referred to as the "active 35 compounds of the invention") may be administered to a patient by various methods, for example, orally as capsules or tablets, parentally as a sterile solution or suspension,

-37-

and in some cases intravenously in the form of a solution. The free base compounds of the invention may be formulated and administered in the form of their pharmaceutically acceptable acid addition salts.

5 The daily dose of the active compounds of the invention is generally in the range of from about 1 to 300 mg/day for the average adult human, and may be administered in single or divided doses.

When incorporated for parenteral administration into a
10 solution or suspension, the active compounds of the invention are present in a concentration of at least 1 weight percent, and preferably between about 4 to 70 weight percent (based on the total weight of the unit). The parenteral dosage unit typically contains between about 5 to
15 100 mg of active compound(s).

The active compounds of the invention may be administered orally with an inert diluent or an edible carrier, or they may be enclosed in gelatin capsules or compressed into tablets. Such preparations should contain
20 at least 0.5% of active compound(s), but the concentration may vary depending upon the particular form and may be from 4 to 70 weight percent (based on the total weight of the unit). The oral dosage unit typically contains between 1.0 mg to 300 mg of active compound.

25 The cholinesterase inhibiting activity of the active compounds of the invention may be determined by a number of standard biological or pharmacological tests. One such procedure for determining cholinesterase inhibition is described by Ellman et al. in "A New and Rapid Colorimetric
30 Determination of Acetylcholinesterase Activity", Biochem. Pharm. 1, 88, (1961).

The present invention is illustrated by the following examples. It will be understood, however, that the invention is not limited to the specific details of these examples.
35 Melting points are uncorrected. Proton nuclear magnetic resonance spectra (^1H NMR) and C^{13} nuclear magnetic resonance spectra (C^{13} NMR) were measured for solutions in


-38-

deuteriochloroform (CDCl_3) except where otherwise noted and peak positions are expressed in parts per million (ppm). The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad.

5 Frequencies (J) are expressed in Hertz. 1 M solutions of lithium diisopropylamide were freshly prepared by adding n-butyllithium (1.6-2.5 M in hexanes) to a solution of diisopropylamine in tetrahydrofuran at 0° .

Example 1

10 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,
2-benzisoxazole maleate



a) 1,4-Piperidinedicarboxylic acid, 1-(1,1-
dimethylethyl)ester, 4-ethyl ester.

A solution of ethyl isonipecotatate (20.0 g, 0.127 mol) and triethylamine (17.8 mL, 0.127 mol) in 1:1 dioxane- H_2O (1.2 L) was cooled to 0°C . After 15 min, t-BOC-anhydride (35.2 g, 0.161 mol) was added and the resulting mixture was allowed to warm to room temperature overnight. The mixture was extracted with ethyl acetate (4 times) and the combined organic layer was washed with 1N hydrochloric acid, water and brine, and then dried (magnesium sulfate), filtered, and concentrated to give a light orange oil. A Kugelrohr distillation (0.05 torr, $80-90^\circ\text{C}$) gave the title carbamate (30.69 g, 94%) as a colorless oil.

20

25 $^1\text{H-NMR}$ (CDCl_3) δ 4.11 (q, 2H, $J=7.2\text{Hz}$), 3.97-4.05 (m, 2H), 2.80 (br t, 2H, $J=11.6\text{Hz}$), 2.40 (tt, 1H, $J=11.0\text{Hz}$, $J=3.9\text{Hz}$), 1.81-1.86 (m, 2H), 1.52-1.66 (m, 2H), 1.43 (s, 9H), 1.23 (t, 3H, $J=7.2\text{Hz}$).

-39-

b) 4-Hydroxymethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester.

Lithium aluminum hydride (4.3 g, 0.114 mol) was added to a cold solution (0°C) of the carbamate formed in step a (26.57 g, 0.103 mol) in tetrahydrofuran (THF) (1L). After 30 min, the ice bath was removed and the reaction mixture was allowed to stir overnight at room temperature. Sodium sulfate decahydrate was added carefully until the evolution of gas subsided. After stirring for 1 hour, the mixture was filtered through a Celite pad and the filtrate was concentrated. Recrystallization (ethyl ether/hexanes) gave the title alcohol (20.67 g, 93%) as a white solid.

¹H-NMR (CDCl₃) δ 4.04-4.26 (m, 2H), 3.49 (d, 2H, J=6.4 Hz), 2.70 (brt, 2H, J=12.0Hz), 1.6-1.73 (m, 3H), 1.47 (s, 9H), 1.15 (ddd, 2H, J=23.2Hz, J=12.0Hz, J=4.3Hz).

c) 4-Iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester.

Triphenylphosphine (31.0 g, 0.119 mol) was added to a mixture of iodine (29.0 g, 0.114 mol) in benzene (1L). After 5 min, pyridine (18.5 ml, 0.228 mol) followed by the alcohol formed in step b (20.5 g, 0.095 mol) was added. The resulting mixture was heated to reflux for 1.5 hours. The cooled reaction mixture was filtered, and the filtrate was washed with saturated sodium thiosulfate (Na₂S₂O₃) and brine, and dried (magnesium sulfate), filtered, and concentrated. Purification by silica gel chromatography (10% → 20% ethyl acetate/hexanes) gave the title iodide (28.5 g, 92%) as a clear oil. Upon cooling, a white solid was obtained.

M.p.: 58-59°C.

¹H-NMR (CDCl₃) δ 4.09 (br d, 2H, J=13.1Hz), 3.08 (d, 2H, J=6.5Hz), 2.66 (br t, 2H, J=13.1Hz), 1.80 (br d, 2H, J=12.9Hz), 1.52-1.64 (m, 1H), 1.43 (s, 9H), 1.11 (ddd, 2H, J=24.7Hz, J=12.7Hz, J=4.3Hz).

d) 4-[2-[1,2-Benzisoxazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester.

A mixture of 3-methyl-1,2-benzisoxazole (0.410 g, 3.08 mmol) and the iodide formed in step c (1.05 g, 3.23 mmol) in

-40-

dry THF (3.2 mL) was cooled to -78°C . Freshly prepared 1M lithium diisopropylamide (LDA) (3.1 mL, 3.1 mmol) was added dropwise and the resulting yellow-orange solution was stirred for 25 min at -78°C . Saturated ammonium chloride was added and the mixture was extracted with ethyl acetate (3 times). The combined organic layer was washed with brine and then dried (MgSO_4), filtered, and concentrated. Purification by silica gel flash chromatography (10% \rightarrow 20% ethyl acetate/hexanes) gave the title compound (0.430 g, 42%) as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3) δ 7.62 (d, 1H, $J=8.0\text{Hz}$), 7.49-7.55 (m, 2H), 7.25-7.31 (m, 1H), 4.09 (m, 2H), 3.00 (t, 2H, $J=7.8\text{Hz}$), 2.66 (br t, 2H, $J=13.0\text{Hz}$), 1.71-1.84 (m, 4H), 1.47-1.53 (m, 1H), 1.43 (s, 9H), 1.14 (ddd, 2H, $J=24.5\text{Hz}$, $J=12.1\text{Hz}$, $J=4.1\text{Hz}$).

e) 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate.

Trifluoroacetic acid (TFA) (7 mL) was added dropwise to a cold (0°C) solution of the piperidine formed in step d (0.50 g, 1.51 mmol) in methylene chloride (7 mL). The resulting solution was stirred at 0°C for 30 min. Volatiles were removed under reduced pressure and excess TFA was removed by concentrating from toluene twice. The crude product was redissolved in methylene chloride (10 mL) and then triethylamine (0.42 mL, 3.01 mmol) and benzyl bromide (0.18 mL, 1.51 mmol) were added. The resulting mixture was stirred overnight (15 hours) at room temperature. The mixture was washed with water and brine and dried (MgSO_4), filtered, and concentrated. Purification by silica gel flash chromatography (50% ethyl acetate/hexanes) gave the title compound (free base) (0.350 g, 73%) as a colorless oil.

The maleate salt was prepared by adding a solution of maleic acid (0.108 g, 0.930 mmol) in ethyl ether (20 mL) to a solution of the free base (0.297 g, 0.926 mmol) in ethyl ether (20 mL). The white solid formed was collected and rinsed with ethyl ether. Yield: 0.35 g, 87%.

-41-

M.p. 146.4-147.6°C.

EIMS (no parent) 319.1, 303.1, 185.2, 172.1, 91.1

¹H NMR (CDCl₃) δ 7.60 (d, J=8, 1H), 7.51-7.52 (m, 2H),
7.37-7.49 (m, 5H), 7.25-7.32 (m, 1H), 6.30 (s, 2H), 4.16 (s,
5 2H), 3.45-3.51 (m, 2H), 2.98 (t, J=7.4, 2H), 2.60-2.70 (m,
2H), 1.84-1.95 (m, 4H), 1.60-1.82 (m, 3H).

¹³C NMR δ 169.5, 163.0, 157.7, 135.8, 131.1, 130.1,
130.0, 129.3, 128.5, 123.5, 121.3, 121.1, 110.0, 60.6, 52.1,
32.8, 28.8, 22.1.


10 IR (KBr) 2944, 2927, 2921, 2499-2518 (broad), 2329-2388
(broad), 1583, 1517, 1473, 1458, 1445, 1438, 1383, 1360, 782
cm⁻¹.

Calc'd for C₂₁H₂₄N₂O⁺C₄H₄O₄:C, 68.79; H, 6.47; N, 6.42.
Found: C, 68.80; H, 6.35; N, 6.27.

15

Example 2

5-Methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-
1,2-benzisoxazole maleate



a) 4-[2-[5-Methyl-1,2-benzisoxazol-3-yl]ethyl]-1-
piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

20 The procedure described in Example 1d was followed
using 3,5-dimethyl-1,2-benzisoxazole (0.500 g, 3.40 mmol),
4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-
dimethylethyl)ester (1.20 g, 3.74 mmol), and 1M lithium
diisopropylamide (LDA) (3.74 mL, 3.74 mmol) in dry THF (5
25 mL). After purification, the title compound (0.910 g, 78%)
was obtained as a clear oil.

¹H-NMR (CDCl₃) δ 7.28-7.40 (m, 3H), 4.04-4.11 (m, 2H),
2.94 (t, 2H, J=7.8Hz), 2.64 (br t, 2H, J=12.3Hz), 2.43 (s,

-42-

3H), 1.70-1.99 (m, 4H), 1.42 (s, 9H), 1.41-1.55 (m, 1H), 1.13 (ddd, 2H, J=24.4Hz, J=12.0Hz, J=4.1Hz).

b) 5-Methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate

5 The procedure described in Example 1e was followed using the piperidine formed in step a (0.910 g, 2.64 mmol) and TFA (13 mL) in methylene chloride (CH₂Cl₂) (13 mL), and triethylamine (3.7 mL, 26.4 mmol) and benzyl bromide (0.32 mL, 2.69 mmol) in methylene chloride (CH₂Cl₂) (20 mL). After
10 purification, the title compound (free base) (0.56 g, 63%) was obtained as a clear oil.

The maleate salt was prepared by adding a solution of maleic acid (0.20 g, 1.72 mmol) in ethyl ether (Et₂O) (10 mL) to a solution of the free base (0.56 g, 1.67 mmol) in Et₂O
15 (40 mL). The white solid formed was collected and rinsed with Et₂O. Yield: 0.70 g, 93%.

M.p. 149-151°C.

¹H NMR (CDCl₃) δ 7.27-7.43 (m, 8H), 6.32 (s, 2H), 4.17 (s, 2H), 3.51 (br d, J=11.6, 2H), 2.96 (t, J=7.3, 2H), 2.66
20 (br t, J=10.8, 2H), 2.45 (s, 3H), 1.60-1.97 (m, 7H).


¹³C NMR δ 169.4, 161.6, 157.3, 135.7, 133.2, 131.6, 131.0, 130.1, 129.3, 128.4, 121.4, 120.2, 60.6, 52.0, 32.8, 32.7, 28.7, 22.0, 21.1.

IR (KBr) 2934, 2848, 2499, 2362, 1701, 1617, 1572,
25 1487, 1454, 1357 cm⁻¹.

EIMS (no parent) 333.1, 317.2, 185.1, 172.1, 91.1 (base).

Calc'd for C₂₂H₂₆N₂O⁺C₄H₄O₄: C, 69.32; H, 6.71; N, 6.22.
Found: C, 69.18; H, 6.48; N, 6.08.

-43-

Example 35,6-Dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate

a) 4-[2-[5,6-Dimethyl-1,2-benzisoxazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

The procedure described in Example 1d was followed using 3,5,6-trimethyl-1,2-benzisoxazole (0.600 g, 3.73 mmol), 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (1.30 g, 4.10 mmol), and 1M LDA (4.10 mL, 4.10 mmol) in dry THF (10 mL). After purification, the title compound (1.04 g, 78%) was obtained as a clear oil.

¹H-NMR (CDCl₃) δ 7.32 (s, 1H), 7.27 (s, 1H), 4.04-4.10 (m, 2H), 2.93 (t, 2H, J=7.8Hz), 2.64 (br t, 2H, J=11.9Hz), 2.35 (s, 3H), 2.32 (s, 3H), 1.70-1.80 (m, 4H), 1.43 (s, 9H), 1.43-1.51 (m, 1H), 1.13 (ddd, 2H, J=24.3Hz, J=12.3Hz, J=4.2Hz).

b) 5,6-Dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate

The procedure described in Example 1e was followed using the piperidine formed in step a (1.04 g, 2.90 mmol) and TFA (16 mL) in CH₂Cl₂ (16 mL), and triethylamine (4.2 mL, 29.0 mmol) and benzyl bromide (0.36 mL, 3.03 mmol) in CH₂Cl₂ (20 mL). After purification, the title compound (free base) (0.53 g, 52%) was obtained as a clear oil.

The maleate salt was prepared by adding a solution of maleic acid (0.18 g, 1.55 mmol) in Et₂O (10 mL) to a solution of the free base (0.53 g, 1.52 mmol) in Et₂O (25 mL). The white solid formed was collected and rinsed with Et₂O. Yield: 0.65 g, 92%.

M.p. 182-183.5 °C.

-44-

^1H NMR (CDCl_3) δ 7.30-7.41 (m, 7H), 6.32 (s, 2H), 4.17 (s, 2H), 3.51 (br d, $J=11.8$, 2H) 2.95 (t, $J=7.2$, 2H) 2.65 (br t, $J=11.7$, 2H), 2.38 (s, 3H), 2.34 (s, 3H), 1.59-1.96 (m, 7H).

5 ^{13}C NMR δ 169.4, 162.3, 157.1, 140.2, 135.7, 132.6, 131.1, 130.1, 129.3, 128.3, 120.3, 119.2, 110.0, 60.6, 52.0, 32.7, 28.7, 22.0, 20.9, 19.9.


EIMS 347.2, 331.1, 185.1, 172.1, 91.1, (base)

10 IR (KBr) 2949, 2914, 2512, 2420, 1580, 1476, 1456, 1449, 1358 cm^{-1} .

Calc'd for $\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}^\circ\text{C}_4\text{H}_4\text{O}_4^\circ 1/4\text{H}_2\text{O}^\circ$: C, 69.14; H, 6.93; N, 5.97. Found: C, 69.27; H, 6.83; N, 5.91.

Example 4

15 5-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinylethyl]-1,2-benzisoxazole maleate



a) 4-[2-[5-Methoxy-1,2-benzisoxazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

The procedure described in Example 1d was followed using 5-methoxy-3-methyl-1,2-benzisoxazole (0.32 g, 1.96 mmol), 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (0.70 g, 2.15 mmol), and 1M LDA (2.0 mL, 2.0 mmol) in dry THF (2 mL). After purification, the title compound (0.62 g, 87%) was obtained as a clear oil.

25 ^1H -NMR (CDCl_3) δ 7.46 (d, 1H, $J=9.1\text{Hz}$), 7.17 (dd, 1H, $J=9.1\text{Hz}$, $J=2.5\text{Hz}$), 6.96 (d, 1H, $J=2.4\text{Hz}$), 4.09-4.16 (m, 2H), 3.87 (s, 3H), 2.99 (t, 2H, $J=7.8\text{ Hz}$), 2.69 (br t, 2H, $J=12.3\text{Hz}$), 1.74-1.85 (m, 4H), 1.46-1.64 (m, 1H), 1.46 (s, 9H), 1.17 (ddd, 2H, $J=22.3\text{ Hz}$, $J=12.2\text{Hz}$, $J=4.2\text{Hz}$).

-45-

b) 5-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate

The procedure described in Example 1e was followed using the piperidine formed in step a (0.58 g, 1.61 mmol) and TFA (7 mL) in CH₂Cl₂ (7 mL), and triethylamine (0.50 mL, 3.6 mmol) and benzyl bromide (0.195 mL, 1.64 mmol) in CH₂Cl₂ (10 mL). After purification, the title compound, free base (0.27 g, 48%) was obtained as a clear oil.

The maleate salt was prepared by adding a solution of maleic acid (0.080 g, 0.69 mmol) in Et₂O (10 mL) to a solution of the free base (0.24 g, 0.68 mmol) in Et₂O (20 mL). The white solid formed was collected and rinsed with Et₂O. Yield: 0.29 g, 91%.

M.p. 143.5-145°C

¹H NMR (CDCl₃) δ 7.35-7.42 (m, 6H), 7.13 (dd, J₁=9.1, J₂=2.5, 1H), 6.92 (d, J=2.4, 1H), 6.30 (s, 2H), 4.17 (s, 2H), 3.83 (s, 3H), 3.46-3.51 (m, 2H), 2.94 (t, J=7.3, 2H), 2.60-2.80 (m, 2H), 1.60-1.96 (m, 7H).

¹³C NMR δ 169.4, 158.4, 157.6, 156.3, 135.8, 131.0, 130.0, 129.2, 128.4, 121.5, 120.3, 110.6, 101.1, 60.5, 56.0, 52.0, 32.7, 32.5, 28.7, 22.0.


IR (KBr) 2942, 2927, 2916, 2518, 2366, 1616, 1572, 1544, 1521, 1480, 1454, 1443, 1384, 1357, 1220 cm⁻¹.

EIMS 349.2, 333.2, 318.1, 259.1, 185.1, 172.1, 91.1 (base).

Calc'd for C₂₂H₂₆N₂O₂·C₄H₄O₄: C, 66.94; H, 6.48; N, 6.00. Found: C, 67.21; H, 6.52; N, 5.94.

Example 5

6-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole



-46-

a) 4-[2-[6-Methoxy-1,2-benzisoxazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

The procedure described in Example 1d was followed using 6-methoxy-3-methyl-1,2-benzisoxazole (0.32 g, 1.96 mmol), 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (0.70 g, 2.15 mmol), and 1M LDA (2.0 mL, 2.0 mmol) in dry THF (3 mL). After purification, the title compound (0.57 g, 80%) was obtained as a white solid.

M.p.: 95-96°C

¹H-NMR (CDCl₃) δ 7.47 (d, 1H, J=8.7Hz), 6.99 (d, 1H, J=2.1Hz) 6.91 (dd, 1H, J=8.6Hz, J=2.1Hz), 4.08-4.11 (m, 2H), 3.89 (s, 3H), 2.97 (t, 2H, J=7.8 Hz), 2.68 (br t, 2H, J=12.7Hz), 1.72-1.84 (m, 4H), 1.46-1.60 (m, 1H), 1.46 (s, 9H), 1.16 (ddd, 2H, J=24.6 Hz, J=12.3Hz, J=4.3Hz).

b) 6-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole

The procedure described in Example 1e was followed using the piperidine formed in step a (0.49 g, 1.36 mmol) and TFA (7 mL) in CH₂Cl₂ (7 mL), and triethylamine (0.85 mL, 6.1 mmol) and benzyl bromide (0.165 mL, 1.39 mmol) in CH₂Cl₂ (8 mL). After purification, the title compound (0.265 g, 55%) was obtained as a white solid.

M.p. 90.5-91.5°C.

¹H NMR (CDCl₃) δ 7.47 (d, J=8.7, 1H), 7.21-7.33 (m, 5H), 6.98 (d, J=1.8, 1H), 6.90 (dd, J₁=8.7, J₂=2.0, 1H), 3.88 (s, 3H), 3.50 (s, 2H), 2.88-2.98 (m, 4H), 1.96 (br t, J=10.6, 2H), 1.74-1.83 (m, 4H), 1.27-1.34 (m, 3H).


¹³C NMR δ 164.8, 162.0, 158.3, 138.4, 129.2, 128.1, 126.9, 121.4, 115.0, 113.8, 92.6, 63.4, 55.7, 53.7, 35.3, 34.3, 32.6, 22.6.

IR (KBr) 2924, 2913, 2797, 2758, 1625, 1608, 1276, 1196, 1154, 734 cm⁻¹.

EIMS 349.2, 333.6, 259.1, 185.1, 172.1, 91.0 (base).

Calc'd for C₂₂H₂₆N₂O₂: C, 75.40; H, 7.48; N, 7.99. Found: C, 75.52; H, 7.63; N, 7.94.

-47-

Example 67-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole fumarate

a) 4-[2-[7-Methoxy-1,2-benzisoxazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

The procedure described in Example 1d was followed using 7-methoxy-3-methyl-1,2-benzisoxazole (0.30 g, 1.84 mmol), 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (0.60 g, 1.85 mmol), and 1M LDA (1.9 mL, 1.9 mmol) in dry THF (2 mL). After purification, the title compound (0.41 g, 62%) was obtained as a pale yellow oil.

¹H-NMR (CDCl₃) δ 7.12-7.19 (m, 2H), 6.91 (dd, 1H, J=6.5Hz, J=2.2Hz), 3.98-4.07 (m, 2H), 3.98 (s, 3H), 2.95 (t, 2H, J=7.8 Hz), 2.62 (br t, 2H, J=12.2Hz), 1.67-1.78 (m, 4H), 1.40-1.48 (m, 1H), 1.40 (s, 9H), 1.10 (ddd, 2H, J=24.5 Hz, J=12.5Hz, J=4.3Hz).

b) 7-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole fumarate

The procedure described in Example 1e was followed using the piperidine formed in step a (0.40 g, 1.11 mmol) and TFA (6 mL) in CH₂Cl₂ (6 mL), and triethylamine (0.34 mL, 2.44 mmol) and benzyl bromide (0.14 mL, 1.18 mmol) in CH₂Cl₂ (10 mL). After purification, the title compound (free base) (0.080 g, 21%) was obtained as a clear oil.

The fumarate salt was prepared by adding a solution of fumaric acid (0.025 g, 0.213 mmol) in ethanol (EtOH) (2 mL) to a solution of the free base (0.071 g, 0.203 mmol) in Et₂O (10 mL). Upon concentration to 4-5 mL, a white/pink solid precipitated. This solid was collected and rinsed with Et₂O. Yield: 0.065 g, 69%.

M.p. 138-139°C.

-48-

^1H NMR (CDCl_3) δ 7.27-7.41 (m, 7H), 7.19 (d, $J=7.7$, 1H), 6.59 (s, 2H), 3.97 (s, 3H), 3.65 (s, 2H), 2.90-3.01 (m, 4H), 2.18 (br t, $J=10.8$, 2H), 1.67-1.77 (m, 4H), 1.26-1.32 (m, 3H).

5 ^{13}C NMR δ 166.6, 158.8, 152.4, 144.0, 136.5, 134.4, 129.4, 128.3, 127.5, 124.9, 122.9, 113.2, 111.4, 61.5, 56.2, 52.6, 34.3, 33.4, 30.7, 22.0.


HRMS Calc'd (free base) 350.1992. Found: 350.1984.

IR (KBr) 1705, 1531, 1266, 756, 642 cm^{-1} .

10 Calc'd for $\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_2 \cdot \text{C}_4\text{H}_4\text{O}_4$: C, 66.94; H, 6.48; N, 6.00. Found: C, 66.76; H, 6.31; N, 5.61.

Example 7

6-Acetamido-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole hemifumarate



15 a) 4-[2-[6-Acetamido-1,2-benzisoxazol-3-yl]ethyl]-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

Freshly prepared 1M LDA (11.0 mL, 11.0 mmol) was added dropwise and quickly to a cold (-78°C) solution of 6-N-acetyl-3-methylbenzisoxazole (1.0 g, 5.26 mol) in THF (50 mL). Immediately after addition was complete, a solution of 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (1.71 g, 5.26 mmol) in THF (8 mL) was added all at once. The resulting yellow-orange solution was stirred for 30 minutes at -78°C . Saturated ammonium chloride (NH_4Cl) was added and the mixture was extracted with ethyl acetate (EtOAc) (3 times). The combined organic layer was washed with brine, dried over magnesium sulfate (MgSO_4), filtered, and concentrated.

Purification by silica gel flash chromatography (20% \rightarrow 30 50% EtOAc/ CH_2Cl_2) gave the title compound (1.56 g, 76%) as a white solid.

-49-

M.p.: 142-143°C

¹H-NMR (CDCl₃) δ 8.76 (s, 1H), 8.05 (s, 1H), 7.48 (d, 1H, J=8.5Hz), 7.32 (dd, 1H, J=8.6Hz, J=1.5Hz), 4.06 (br d, 2H, J=11.5Hz), 2.94 (t, 2H, J=7.8Hz), 2.66 (br t, 2H, J=11.8 Hz), 2.20 (s, 3H), 1.69-1.80 (m, 4H), 1.41-1.47 (m, 1H), 1.44 (s, 9H), 1.12 (ddd, 2H, J=23.8 Hz, J=12.0Hz, J=3.9Hz).

b) 6-Acetamido-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole fumarate

Trifluoroacetic acid (TFA) (4 mL) was added dropwise to a cold (0°C) solution of the piperidine formed in step a (0.40 g, 1.03 mmol) in CH₂Cl₂ (8 mL). The resulting solution was stirred at 0°C for 30 min. Volatiles were removed under reduced pressure and excess TFA was removed by concentrating from toluene twice. The crude product was redissolved in CH₂Cl₂ (10 mL) and triethylamine (1.44 mL, 10.3 mmol) and benzyl bromide (0.184 mL, 1.55 mmol) was added. The resulting mixture was stirred for 6 hours at room temperature. The mixture was washed with water and brine and dried (MgSO₄), filtered, and concentrated. Purification by silica gel flash chromatography (CH₂Cl₂ → 10% MeOH/CH₂Cl₂) gave the title compound (free base) (0.270 g, 69%) as a white solid.


The fumarate salt was prepared by adding a solution of fumaric acid (0.091 g, 0.788 mmol) in ethanol (EtOH) (5 mL) to a solution of the free base (0.270 g, 0.716 mmol) in CH₂Cl₂ (20 mL). After concentrating, the solid obtained was recrystallized from EtOH to give white needles. Yield: 0.17 g, 48%.

M.p. 225-226°C.

¹H NMR (DMSO-d₆) δ 10.37 (s, 1H), 8.13 (s, 1H), 7.76 (d, 1H, J=8.5), 7.25-7.36 (m, 6H), 6.59 (s, 2H), 3.54 (s, 2H), 2.83-2.96 (m, 4H), 2.10 (s, 3H), 2.01 (br t, 2H, J=11.1), 1.69-1.73 (m, 4H), 1.20-1.28 (m, 3H).

Calc'd for C₂₃H₂₇N₃O₂•1/2C₄H₄O₄•1/4H₂O: C, 68.24; H, 6.76; N, 9.55. Found: C, 68.35; H, 6.63; N, 9.35.

-50-

Example 86-Amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate

A mixture of 6-acetamido-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole (0.30 g, 0.79 mmol) in 1N HCl (10 mL) was heated to reflux for 30 min. The cooled reaction mixture was made basic with 10% NaOH and extracted with EtOAc (2 times). The combined organic layer was washed with brine and dried (MgSO₄), filtered, and concentrated to give the title compound (free base) (0.259 g, 98%) as an oil.

The mono maleate salt was prepared by adding a solution of maleic acid (0.099 g, 0.85 mmol) in EtOH (5 mL) to a solution of the free base (0.26, 0.77 mmol) in CH₂Cl₂ (3 mL). After concentrating, the residue was triturated with Et₂O to give a white powder. Yield: 0.29 g, 64%.

M.p. 173.0-173.5°C.

¹H NMR (DMSO-d₆) δ 7.41-7.47 (m, 6H), 6.58-6.63 (m, 2H), 6.06 (s, 2H), 5.87 (br s, 2H), 4.26 (s, 2H), 3.29-3.38 (m, 2H), 2.80-2.95 (m, 4H), 1.90 (br d, J=12.5, 2H) 1.25-1.80 (m, 5H).


¹³C NMR (DMSO-d₆) 167.4, 164.8, 157.4, 151.9, 136.0, 131.2, 130.1, 129.5, 128.9, 121.9, 112.8, 110.7, 90.9, 59.3, 51.5, 32.6, 32.6, 28.6, 21.6.

EIMS (no parent) 289, 268, 218, 190 (base).

IR (KBr) 3483, 3384, 2929, 2526, 1633, 1619, 1582, 1515, 1474, 1459, 1438, 1389, 1379, 1359, 877, 702 cm⁻¹.

Calc'd for C₂₁H₂₅N₂O•C₄H₄O₄:C, 66.50; H, 6.47; N, 9.31. Found: C, 66.49; H, 6.43; N, 9.22.

-51-

Example 96-Benzamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleatea) 6-Benzamide-3-methyl-1,2-benzisoxazole

5 Benzoyl chloride (0.56 mL, 4.82 mmol) was added to a solution of 6-amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole (0.70 g, 4.72 mmol), triethylamine (1.35 mL, 9.69 mmol), and 4-dimethylaminopyridine (0.07 g, 0.57 mmol) in CH₂Cl₂ (30 mL). The resulting mixture was stirred overnight at room temperature. The heterogenous mixture was concentrated, and the solid obtained was collected and washed with water and ether and air-dried to give the title compound (1.02 g, 86%) as an off-white solid. A small sample was purified by recrystallization from EtOH to give pure white flakes.

M.p.: 213-214°C.

¹H-NMR (DMSO-d₆) δ 10.6 (s, 1H), 8.30 (s, 1H), 7.98 (d, 2H, J=6.9Hz), 7.80 (d, 1H, J=8.6Hz), 7.68 (d, 1H, J=8.8Hz), 7.52-7.63 (m, 3H), 2.53 (s, 3H).

b) 4-[2-[6-Benzamide-1,2-benzisoxazole-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

The procedure described in Example 7a was followed using the benzamide formed in step a (1.0 g, 3.96 mmol), 1M LDA (7.95 mL, 7.95 mmol), and 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (1.30 g, 4.00 mmol) in dry THF (50 mL), except that after addition of reagents, the mixture was stirred at -78°C for 1.5 hours. After purification by chromatography (30% → 50% EtOAc/hexanes), the title compound (1.54 g, 87%) was obtained as a pale yellow solid. A small sample was

-52-

purified by recrystallization ($\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$) to give a white solid.

M.p.: 177-178.5°C

^1H -NMR (CDCl_3) δ 8.61 (s, 1H), 8.15 (s, 1H), 7.86 (d, 2H, $J=7.4\text{Hz}$), 7.39-7.53 (m, 5H), 4.03 (br d, 2H, $J=12.7\text{Hz}$), 2.92 (t, 2H, $J=8.0\text{Hz}$), 2.50-2.73 (m, 2H), 1.60-1.80 (m, 4H), 1.40-1.45 (m, 1H), 1.41 (s, 9H), 1.13 (ddd, 2H, $J=24.0\text{ Hz}$, $J=12.2\text{Hz}$, $J=3.8\text{Hz}$).

c) 6-Benzamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole maleate

Trifluoroacetic acid (TFA) (8.4 mL) was added dropwise to a cold (0°C) solution of the piperidine formed in step b (0.70 g, 1.56 mmol) in CH_2Cl_2 (10 mL). The resulting solution was stirred at 0°C for 30 min. Volatiles were removed under reduced pressure and excess TFA was removed by concentrating from toluene twice. The crude product was dissolved in THF (10 mL) and triethylamine (2.1 mL, 15.1 mmol) followed by benzyl bromide (0.21 mL, 1.77 mmol) was added. The resulting mixture was diluted with ethyl acetate and stirred overnight at room temperature. The mixture was then washed with water and brine and dried (MgSO_4), filtered, and concentrated. Purification by silica gel flash chromatography (30% EtOAc/ $\text{CH}_2\text{Cl}_2 \rightarrow 100\%$ EtOAc) gave the title compound (free base) (0.280 g, 41%) as a pale yellow solid.

The maleate salt was prepared by adding a solution of maleic acid (0.081 g, 0.702 mmol) in EtOH (5 mL) to a solution of the free base (0.280 g, 0.638 mmol) in CH_2Cl_2 (20 mL). After concentrating, the solid obtained was recrystallized from EtOH/ CH_2Cl_2 to give a white solid. Yield: 0.208 g, 59%.

M.p. 181.5-183.0°C.

^1H NMR ($\text{DMSO}-d_6$) δ 10.65 (s, 1H), 8.31 (s, 1H), 7.98 (d, $J=7.2$, 2H), 7.85 (d, $J=8.6$, 1H), 7.48-7.71 (m, 9H), 6.03 (s, 2H), 4.25 (br s, 2H), 3.20-3.60 (m, 4H), 2.89-3.02 (t, @ 2.99, $J=7.5$ and m, 4H), 1.40-1.97 (m, 7H).

^{13}C NMR ($\text{DMSO}-d_6$) δ 167.2, 166.2, 163.0, 158.1, 141.5, 135.9, 134.6, 132.0, 131.2, 129.5, 128.9, 128.5, 127.8,

-53-

121.9, 117.01, 116.8, 99.9, 59.5, 51.7, 32.8, 32.5, 28.8, 21.7.


EIMS 439.2, 422.2 (100), 383, 348, 293, 185.

IR (KBr) 2934, 2919, 1657, 1610, 1579, 1536, 1499,
5 1491, 1462, 1453, 1352 cm^{-1} .

Calc'd for $\text{C}_{28}\text{H}_{29}\text{N}_3\text{O}_2 \cdot \text{C}_4\text{H}_4\text{O}_4$: C, 69.17; H, 5.99; N, 7.56.
Found: C, 68.81; H, 5.90; N, 7.49.

Example 10

6-Benzenesulfonamide-3-[2-[1-(phenylmethyl)-4- 10 piperidinyl]ethyl]-1,2-benzisoxazole fumarate



a) 6-Benzenesulfonamide-3-methyl-1,2-benzisoxazole

Benzenesulfonyl chloride (0.528 mL, 4.14 mmol) was added to a cold (0°C) solution of 6-amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole (0.613
15 g, 4.14 mmol) and pyridine (0.670 mL, 8.28 mmol) in CH_2Cl_2 (30 mL). After 1.3 hours, saturated sodium bicarbonate (NaHCO_3) was added and the resulting mixture was stirred overnight at room temperature. The organic layer was separated and washed with water and brine and dried (MgSO_4),
20 filtered, and concentrated. Purification by silica gel flash chromatograph (5% EtOAc/ CH_2Cl_2) gave the title compound (0.867 g, 83%) as a white solid.

M.p.: $183-184^\circ\text{C}$.

$^1\text{H-NMR}$ (CDCl_3) δ 10.9 (br s, 1H), 7.84 (d, 2H, $J=6.7\text{Hz}$),
25 7.68 (d, 1H, $J=8.5\text{Hz}$), 7.52-7.63 (m, 3H), 7.34 (d, 1H, $J=1.5\text{Hz}$), 7.11 (dd, 1H, $J=8.5\text{Hz}$, $J=1.7\text{Hz}$), 2.52 (s, 3H).

b) 4-[2-[6-Benzenesulfonamide-1,2-benzisoxazol-3- yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

30 The procedure described in Example 7a was followed using the benzenesulfonamide formed in step a (0.60 g, 2.08

-54-

mmol), 1M LDA (4.58 mL, 4.58 mmol), and 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1-dimethylethyl) ester (0.813 g, 2.50 mmol) in dry THF (70 mL), except that after addition of reagents, the mixture was stirred at -78°C for 10-15 min.

- 5 After purification by chromatography (20% → 40% EtOAc/hexanes), the title compound (0.997 g, 99%) was obtained as a white foam.

M.p.: 66-67°C.

¹H-NMR (CDCl₃) δ 7.85 (dd, 2H, J=8.3Hz, J=1.6Hz), 7.35-
10 7.57 (m, 6H), 7.02 (dd, 1H, J=8.5Hz, J=1.6Hz), 4.11 (br d, 2H, J=13.2Hz), 2.94 (t, 2H, J=7.4Hz), 2.68 (br t, 2H, J=12.8Hz), 1.71-1.77 (m, 4H), 1.46 (s, 9H), 1.46-1.55 (m, 1H), 1.15 (ddd, 2H, J=23.6Hz, J=11.7Hz, J=3.9Hz).

- c) 6-Benzenesulfonamide-3-[2-[1-(phenylmethyl)-4-
15 piperidinyl]ethyl]-1,2-benzisoxazole fumarate

Trimethylsilyl trifluoromethanesulfonate (1.30 mL, 6.76 mmol) was added dropwise to a cold (0°C) solution of the piperidine formed in step b (0.819 g, 1.69 mmol) and 2,6-lutidine (0.590 mL, 5.07 mmol) in CH₂Cl₂ (17 mL). After 1.5
20 hours, saturated sodium bicarbonate (NaHCO₃) was added and the resulting mixture was stirred at room temperature for 15 min. The white precipitate formed was collected by filtration and redissolved in water at pH 2. This acidic aqueous layer was extracted with CH₂Cl₂ (2 times) and EtOAc
25 (1 time). All organic layers were combined, dried (MgSO₄), filtered, and concentrated. The crude white solid obtained was suspended in THF (30 mL) and DMF (50 mL), and triethylamine (0.40 mL, 2.86 mmol) and benzyl bromide (0.22 mL, 1.86 mmol) were added. The resulting heterogenous
30 mixture was stirred at room temperature for 24 hours (with time a more homogenous mixture was obtained). The mixture was concentrated, and CH₂Cl₂ was added to the residue. The organic layer was washed with water and brine and dried (MgSO₄), filtered, and concentrated. Purification by silica
35 gel flash chromatography (CH₂Cl₂ → 5% MeOH/CH₂Cl₂) gave the title compound (free base) (0.334 g, 49%) as a white foam.


-55-

The fumarate salt was prepared by adding a solution of fumaric acid (0.040 g, 0.345 mmol) in EtOH (3 mL) to a solution of the free base (0.150 g, 0.315 mmol) in CH₂Cl₂ (6 mL). After concentrating, the residue was triturated with Et₂O to give a white solid. Yield: 0.151 g, 81%.

¹H NMR (DMSO-d₆) δ 7.82 (d, 2H, J=7.0), 7.70 (d, 1H, J=8.6), 7.51-7.60 (m, 3H), 7.27-7.32 (m, 6H), 7.07 (dd, 1H, J₁=8.6, J₂=1.6), 6.59 (s, 2H), 3.55 (s, 2H), 2.82-2.90 (m, 4H), 2.03 (br t, 2H, J=11.5), 1.61-1.70 (m, 4H), 1.20-1.25 (m, 3H).

Example 11

6-(4-Morpholinyl)-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole



a) 3-Methyl-6-(4-morpholinyl)-1,2-benzisoxazole

A mixture of 6-amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole (0.230 g, 1.55 mmol), β, β'-dibromodiethyl ether (0.397 g, 1.71 mmol), and diisopropylethyl amine (Hunig's base, 0.648 mL, 3.72 mmol) in toluene (2.5 mL) was heated at 120°C for 15 hours. The cooled reaction mixture was diluted with EtOAc and washed with water and brine and dried (MgSO₄), filtered, and concentrated. Two additional separate reactions using 6-amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole (0.050 g, 0.34 mmol, and 0.150 g, 1.01 mmol) were carried out in the same manner. Crude product from the three reactions was combined and purified by silica gel flash chromatography (1% MeOH/CH₂Cl₂) to give the title compound (0.499 g, 79% combined yield) as a pale yellow solid. A small sample was further purified by recrystallization (EtOAc/hexanes) to give a white solid.

-56-

M.p.: 138.5-139.5°C.

¹H-NMR (CDCl₃) δ 7.46 (d, 1H, J=8.7Hz), 6.95 (dd, 1H, J=8.7Hz, J=2.0Hz), 6.90 (d, 1H, J=1.9Hz), 3.88 (t, 4H, J=4.9Hz), 3.27 (t, 4H, J=4.8Hz), 2.52 (s, 3H).

5 b) 4-[2-[6-(4-Morpholinyl)-1,2-benzisoxazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

The procedure described in Example 7a was followed using the morpholino derivative formed in step a (0.369 g, 1.69 mmol), 1M LDA (1.86 mL, 1.86 mmol), and 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (0.605 g, 1.86 mmol) in dry THF (8 mL). Another reaction with the above morpholino derivative (0.100 g, 0.46 mmol) was also carried out. Crude product from both reactions was combined and after purification by chromatography (5% → 40% EtOAc/hexanes), the title compound (0.477 g, 53%) was obtained as a white solid. A small sample was further purified by recrystallization (EtOAc/hexanes) to give a white solid.

20 M.p.: 164-165°C.

¹H-NMR (CDCl₃) δ 7.46 (d, 1H, J=8.6Hz), 6.92-6.97 (m, 2H), 4.02-4.15 (m, 2H), 3.89 (t, 4H, J=4.9Hz), 3.27 (t, 4H, J=4.9Hz), 2.95 (t, 2H, J=7.8Hz), 2.7 (br t, 2H, J=12.1Hz), 1.74-1.80 (m, 4H), 1.46-1.56 (m, 1H), 1.46 (s, 9H), 1.10-1.22 (m, 2H).

25 c) 6-(4-Morpholinyl)-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole

Trifluoroacetic acid (TFA) (5 mL) was added to a cold (0°C) solution of the piperidine formed in step b (0.40 g, 0.96 mmol) and thioanisole (1.13 mL, 9.60 mmol) in CH₂Cl₂ (10 mL). The resulting mixture was stirred at 0°C for 30 min. Volatiles were removed under reduced pressure and 1N sodium hydroxide (NaOH) was added to the residue. The aqueous layer was extracted with EtOAc (2 times) and the combined organic layer was washed with water and brine and dried (MgSO₄), filtered, and concentrated. The yellow oil obtained was redissolved in CH₂Cl₂ (10 mL), and triethylamine (0.267

-57-

mL, 1.92 mmol) and benzyl bromide (0.148 mL, 1.25 mmol) were added. The resulting mixture was stirred overnight at room temperature. The mixture was washed with water and brine and dried (MgSO_4), filtered, and concentrated. Purification
 5 by silica gel flash chromatography ($\text{CH}_2\text{Cl}_2 \rightarrow 10\% \text{ MeOH}/\text{CH}_2\text{Cl}_2$) gave the title compound (0.285 g, 73%) as a white solid. A small sample was further purified by recrystallization (EtOH) to give a white solid.

M.p. 129-130°C.

10 ^1H NMR (CDCl_3) δ 7.44 (d, $J=8.7$, 1H), 7.23-7.30 (m, 5H), 6.89-6.93 (m, 2H), 3.87 (t, $J=4.8$, 4H), 3.47 (s, 2H), 3.25 (t, $J=4.8$, 4H), 2.85-2.94 (m, 4H), 1.92 (br t, $J=10.8$, 2H), 1.72-1.79 (m, 4H), 1.21-1.31 (m, 3H).

^{13}C NMR 165, 158, 153.4, 138.8, 129.2, 128.1, 126.9,
 15 121.3, 113.3, 94.5, 66.7, 63.5, 53.7, 49.1, 35.3, 34.4, 32.1, 22.7.

IR (KBr) 3020, 2920, 1950, 1895, 1815, 1722, 1620, 1450, 1250, 1122, 738 cm^{-1} .

Calc'd for $\text{C}_{25}\text{H}_{31}\text{N}_3\text{O}_2$: C, 74.04; H, 7.70; N, 10.36.


20 Found: C, 73.75; H, 7.69; N, 10.36.

Example 12

5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-
6H-

pyrrolo [4,5-f]-1,2-benzisoxazol-6-one maleate

25



30

a) 5-Acetyl-1,3-dihydro-6-hydroxy-2H-indol-2-one

Acetyl chloride (4.09 mL, 0.0575 mol) was added to a
 35 slurry of aluminum trichloride (AlCl_3) (35.36 g, 0.265 mol) in carbon disulfide (CS_2) (250 mL). After 2-3 min, 6-

-58-

methoxyoxindole (7.22 g, 0.0442 mol) was added. The resulting mixture was heated to reflux for 2.5 hours. (A black tar developed with time). Excess solvent was decanted and ice water was added carefully to the residue. The resulting mixture was stirred overnight. The pale yellow solid obtained was collected, washed with water and dried under high vacuum to give the title compound (7.32 g, 87%).

¹H-NMR (DMSO-d₆) δ 13.0 (s, 1H), 10.8 (s, 1H), 7.70 (s, 1H), 6.30 (s, 1H), 3.40 (s, 2H), 2.54 (s, 3H).

10 b) 5-Acetyl-1,3-dihydro-6-hydroxy-2H-indol-2-one, 5-oxime

An aqueous solution of hydroxylamine hydrochloride (8.26 g, 0.119 mol) and sodium acetate trihydrate (16.9 g, 0.124 mol) was added to a mixture of the ketone formed in step a (9.88 g, 0.0517 mol) in EtOH (600 mL). The resulting mixture was heated to reflux for 20 hours. The hot reaction mixture was filtered and the solid collected was rinsed with EtOH. After drying, the title compound (10.11 g, 95%) was obtained as a pale yellow solid.

20 ¹H-NMR (DMSO-d₆) δ 12.0 (s, 1H), 11.4 (s, 1H), 10.5 (s, 1H), 7.29 (s, 1H), 6.35 (s, 1H), 3.38 (s, 2H), 2.20 (s, 3H).

c) 5-Acetyl-1,3-dihydro-6-hydroxy-2H-indol-2-one, 5-oxime acetate

A heterogenous mixture of the oxime formed in step b (7.15 g, 34.7 mmol) in acetic anhydride (55 mL) was heated at 80°C for 2 hours. The cooled reaction mixture was filtered and the solid collected was rinsed with water. After drying, the title compound (4.67 g, 54%) was obtained as a pale yellow solid.

30 ¹H-NMR (DMSO-d₆) δ 11.3 (s, 1H), 10.6 (s, 1H), 7.35 (s, 1H), 6.44 (s, 1H), 3.41 (s, 2H), 2.37 (s, 3H), 2.21 (s, 3H).

d) 5,7-Dihydro-3-methyl-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one

A mixture of the oxime acetate formed in step c (4.48 g, 18.0 mmol) and pyridine (14.6 mL, 180 mmol) in dimethylformamide (DMF) (660 mL) was heated at 125-130°C for 4 hours. The cooled reaction mixture was poured over water

-59-

and extracted with EtOAc (4 times). The combined organic layer was washed with water and brine and dried (MgSO₄), filtered, and concentrated. Purification by silica gel flash chromatography (50% EtOAc/hexanes → 100% EtOAc) gave the title compound (2.20 g, 65%) as a pale yellow-orange solid.

M.p. (EtOAc): 264-265°C (dec).

¹H-NMR (DMSO-d₆) δ 10.8 (s, 1H), 7.60 (s, 1H), 6.98 (s, 1H), 3.57 (s, 2H), 2.47 (s, 3H).

10 e) 4-[2-[5,7-Dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester.

The procedure described in Example 7a was followed using the benzisoxazole formed in step d (2.33 g, 12.4 mmol), 1M LDA (40.9 mL, 40.9 mmol), and 4-iodomethyl-1-piperidine-carboxylic acid, 1-(1,1-dimethylethyl) ester (4.42 g, 13.6 mmol) in dry THF (500 mL), except that after addition of reagents, the mixture was stirred at -78°C for 4 hours. Purification by chromatography (20% → 30% EtOAc/CH₂Cl₂) gave recovered starting material (0.210 g, 9%) and the title compound (2.75 g, 58%) as an off-white solid.

¹H-NMR (CDCl₃) δ 8.48 (s, 1H), 7.44 (s, 1H), 7.03 (s, 1H), 4.08-4.14 (m, 2H), 3.63 (s, 2H), 2.97 (t, 2H, J=7.8Hz), 2.69 (br t, 2H, J=12.8Hz), 1.74-1.84 (m, 4H), 1.46-1.55 (m, 1H), 1.46 (s, 9H), 1.18 (ddd, 2H, J=24.4Hz, J=12.1Hz, J=4.3Hz).

f) 5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one maleate

30 Trifluoroacetic acid (TFA) (3.3 mL) was added dropwise to a cold (0°C) solution of the piperidine formed in step e (0.50 g, 1.30 mmol) in CH₂Cl₂ (13 mL). After 30 min, the mixture was concentrated and excess TFA was removed by concentrating from toluene (2 to 3 times). The crude residue was dissolved in DMF (12.5 mL) and sodium carbonate (Na₂CO₃) (0.689 g, 6.50 mmol) and benzyl bromide (0.186 mL, 1.56 mmol) were added. The resulting mixture was stirred at

-60-

room temperature for 4 hours. The reaction was filtered and the filtrate was concentrated in vacuo. The residue was dissolved in methylene chloride and washed with brine, and dried (MgSO_4), filtered, and concentrated. Purification by silica gel chromatography ($\text{CH}_2\text{Cl}_2 \rightarrow 10\%$ methanol/ CH_2Cl_2) gave the title compound (free base) (0.343 g, 70%) as a white solid.

The maleate salt was prepared by adding a solution of maleic acid (0.061 g, 0.528 mmol) in ethanol (EtOH) (1 mL) to a solution of the free base (0.180 g, 0.48 mmol) in CH_2Cl_2 (10 mL). After concentrating, the salt was purified by recrystallization from isopropanol to give an off-white solid. Yield: 0.173 g, 73%.


M.p. 194-195°C.

^1H NMR ($\text{DMSO}-d_6$) δ 10.82 (s, 1H), 7.65 (s, 1H), 7.48 (s, 5H), 7.00 (s, 1H), 6.03 (s, 1H), 4.24 (br s, 2H), 3.58 (s, 2H), 3.25-3.38 (m, 2H), 2.94 (t, 2H, $J=7.6$), 2.81-2.97 (m, 2H), 1.86-1.96 (m, 2H), 1.62-1.76 (m, 2H), 1.30-1.60 (m, 3H).

Calc'd for $\text{C}_{23}\text{H}_{25}\text{N}_3\text{O}_2 \cdot \text{C}_4\text{H}_4\text{O}_4$: C, 65.97; H, 5.95; N, 8.55. Found: C, 65.98; H, 6.04; N, 8.54.

Example 13

1-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]isoquinoline maleate



a) 4-[2-[1-Isoquinolyl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester

The procedure described in Example 7a was followed using 1-methyl isoquinoline (0.50 g, 3.49 mmol), 1M LDA (4.2 mL, 4.2 mmol), and 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl) ester (1.2 g, 3.84 mmol) in dry THF (45 mL) except that after addition of the reagents, the

-61-

mixture was stirred at -78°C for 1.75 hours. Purification by chromatography (30% EtOAc/toluene) gave the title compound (0.784 g, 66%) as a yellow oil.

¹H-NMR (CDCl₃) δ 8.38 (d, 1H, J=5.8Hz), 8.08 (d, 1H, J=8.3Hz), 7.77 (d, 1H, J=8.0Hz), 7.62 (t, 1H, J=7.1Hz), 7.54 (t, 1H, J=7.1Hz), 7.46 (d, 1H, J=5.7Hz), 4.08 (br s, 2H), 3.25-3.30 (m, 2H), 2.67 (br t, 2H, J=12.3Hz), 1.73-1.81 (m, 4H), 1.49-1.63 (m, 1H), 1.42 (s, 9H), 1.17 (ddd, 2H, J=24.6Hz, J=12.1Hz, J=3.8Hz).

b) 1-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-isoquinoline maleate


The procedure described in Example 1e was followed using the piperidine formed in step a (0.713 g, 2.10 mmol) and TFA (7 mL) in CH₂Cl₂ (14 mL), and triethylamine (2.9 mL, 21.0 mmol) and benzyl bromide (0.275 mL, 2.31 mmol) in CH₂Cl₂ (60 mL). After purification (CH₂Cl₂ → 5% MeOH/CH₂Cl₂), the title compound (free base) (0.26 g, 38%) was obtained as a pale yellow oil.

The mono maleate salt was prepared by adding a solution of maleic acid (0.10 g, 0.867 mmol) in EtOH (3 mL) to a solution of the free base (0.26 g, 0.788 mmol) in CH₂Cl₂ (7 mL). After concentrating, the salt was purified by recrystallization [cold (0°) EtOAc] to give an off-white solid. Yield: 0.195 g, 56%.

¹H NMR (DMSO-d₆) δ 8.39 (d, 1H, J=5.7), 8.26 (d, 1H, J=8.3), 7.97 (d, 1H, J=8.1), 7.77 (t, 1H, J=7.4), 7.66-7.71 (m, 2H), 7.49 (s, 5H), 6.05 (s, 2H), 4.28 (br s, 2H), 3.27-3.32 (m, 2H), 2.87-2.90 (m, 2H), 1.76-2.03 (m, 6H), 1.55-1.69 (m, 1H), 1.33-1.46 (m, 2H).

Example 14

3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisothiazole maleate



-62-

a) 4-[2-[1,2-Benzisothiazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester.

The procedure described in Example 1d was followed using 3-methyl-1,2-benzisothiazole (0.50 g, 3.35 mmol), 4-
5 iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (1.20 g, 3.69 mmol), and 1M lithium diisopropylamide (LDA) (3.35 mL, 3.35 mmol) in dry THF (100 mL). After purification, the title compound (0.582 g, 50%) was obtained as a pale yellow oil.

10 ¹H-NMR (CDCl₃) δ 7.91-7.96 (m, 2H), 7.43-7.52 (m, 2H), 4.05-4.14 (m, 2H), 3.15 (t, 2H, J=7.9Hz), 2.69 (br t, 2H, J=12.1Hz), 1.74-1.88 (m, 4H), 1.46-1.60 (m, 1H), 1.46 (s, 9H), 1.29-1.10 (m, 2H).

b) 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisothiazole maleate

The procedure described in Example 1e was followed using the piperidine formed in step a (0.102 g, 0.29 mmol) and trifluoroacetic acid (TFA) (0.75 mL) in methylene chloride (CH₂Cl₂) (3 mL), and triethylamine (0.202 mL, 1.45
20 mmol) and benzyl bromide (0.038 mL, 0.32 mmol) in methylene chloride (CH₂Cl₂) (3 mL). After purification, the title compound (free base) (0.058 g, 62%) was obtained as a clear oil.

The maleate salt was prepared by adding a solution of
25 maleic acid (0.020 g, 0.17 mmol) in EtOH (3 mL) to a solution of the free base (0.058 g, 0.17 mmol) in CH₂Cl₂ (3 mL). The resulting mixture was concentrated and the residue was triturated with Et₂O. The white solid obtained was filtered and collected to give the title compound (0.077 g,
30 96%).

¹H-NMR (CDCl₃) δ 8.14-8.21 (m, 2H), 7.63 (t, J=7.4, 1H), 7.50-7.55 (m, 6H), 6.04 (s, 2H), 4.26 (br s, 2H), 3.35 (br s, 2H), 3.15 (t, J=7.6, 2H), 2.80-2.92 (m, 2H), 1.92-2.00 (m, 2H), 1.78-1.88 (m, 2H), 1.54-1.65 (m, 1H), 1.35-1.45 (m,
35 2H).

M.p.: 175-176°C.

-63-

^{13}C -NMR (DMSO- d_6) δ 167.3, 166.2, 151.6, 136.0, 134.2, 131.2, 130.1, 129.5, 128.9, 127.9, 124.9, 123.6, 120.6, 59.4, 51.6, 32.5, 33.5, 28.9, 27.8.


IR (KBr) 3030, 2910, 2350, 1700, 1575, 1445, 1350 cm^{-1} .

5 EIMS: 336.2 (M^+ , free base), 319.1, 245.1, 185.1, 172.1, 91.0 (base).

Anal. Calcd. for $\text{C}_{21}\text{H}_{24}\text{N}_2\text{S}^+\text{C}_4\text{H}_4\text{O}_4$: C, 66.35; H, 6.24; N, 6.19. Found: C, 66.21; H, 5.93; N, 6.03.

Example 15

10 4-[2-[1-(Phenylmethyl)-4-piperidyl]ethyl]-1,3-quinazoline maleate



a) 4-[2-[1,3-Quinazol-4-yl]ethyl]-1-piperidine-carboxylic acid, 1-(1,1-dimethylethyl)ester

20 Freshly made 1M LDA (4.2 mL, 4.2 mmol) was added to a solution of 4-methyl-1,3-quinazoline (0.60 g, 4.2 mmol) in THF (35 mL) at 0°C. To the yellow solution obtained, was added neat trimethylsilyl chloride (0.53 mL, 4.2 mmol). The ice bath was removed and the reaction was allowed to stir
25 for 3 min. The mixture was re-cooled to 0°C, and a second equivalent of 1M LDA (4.2 mL, 4.2 mmol) was added. Next, a solution of 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (1.49 g, 4.6 mmol) in THF (10 mL) was added and stirring was continued at 0°C for 1 hour.
30 Dilute HCl (1 N) was added and the mixture was stirred at room temperature for 30 min. The reaction was made basic by addition of 1N NaOH and extracted with EtOAc. The organic layer was washed with brine, dried (MgSO_4), filtered, and concentrated. Purification by silica gel flash
35 chromatography (20→50% EtOAc-hexanes) gave the title compound (0.466 g, 33%) as an oil.

-64-

¹H-NMR (CDCl₃) δ 9.21 (s, 1H), 8.12 (d, 1H, J=7.7Hz), 8.05 (d, 1H, J=8.4Hz), 7.90 (t, 1H, J=7.8 Hz), 7.65 (t, 1H, J=7.7Hz), 4.08-4.17 (m, 2H), 3.32 (t, 2H, J=8.3Hz), 2.72 (br t, 2H, J=12.1Hz), 1.78-1.90 (m, 4H), 1.57-1.59 (m, 1H), 1.47 (s, 9H), 1.18-1.29 (m, 2H).

b) 4-[2-[1-(Phenylmethyl)-4-piperidyl]ethyl]-1,3-guinazoline maleate

The procedure described in Example 1e was followed with the piperidine obtained in step a (0.429 g, 1.26 mmol) and TFA (3.5 mL) in CH₂Cl₂ (13 mL), and triethylamine (0.88 mL, 6.3 mmol) and benzyl bromide (0.17 mL, 1.39 mmol) in CH₂Cl₂ (22 mL). After purification by chromatography (CH₂Cl₂→10% MeOH-CH₂Cl₂), the title compound, free base (0.179 g, 43%) was obtained as a colorless oil.

The maleate salt was prepared by adding a solution of maleic acid (0.052 g, 0.45 mmol) in Et₂O (10 mL) to a solution of the free base (0.135 g, 0.41 mmol) in Et₂O (200 mL). The white solid obtained was collected by filtration to give the title compound (0.103 g, 56%).

M.p.: 121-122°C.

¹H-NMR (DMSO-d₆) δ 9.16 (s, 1H), 8.33 (d, 1H, J=8.4Hz), 7.98-8.01 (m, 2H), 7.72-7.78 (m, 1H), 7.43 (s, 5H), 6.02 (s, 2H), 4.08 (br s, 2H), 3.32 (t, 2H, J=7.7Hz), 3.15-3.35 (m, 2H), 2.60-2.80 (m, 2H), 1.89-1.94 (m, 2H), 1.76-1.79 (m, 2H), 1.30-1.60 (m, 3H).

IR (KBr) 3037, 2923, 1705, 1571, 1386 cm⁻¹.


EIMS: 331 (M⁺, free base), 318, 248, 143, 77 (base).

Anal. Calcd. for C₂₂H₂₅N₃O₄: C, 69.78; H, 6.53; N, 9.39. Found: C, 69.46; H, 6.61; N, 9.26.

-65-

Example 166-Hydroxy-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole

5



10 A mixture of 6-methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole (0.11 g, 0.288 mmol) in 48% aqueous HBr (10 mL) was heated at 110°C for 16 hours. The mixture was made basic by addition of saturated NaHCO₃ and extracted with CH₂Cl₂. The organic layer was dried
 15 (MgSO₄), filtered, and concentrated. Purification by silica gel flash chromatography (3→6% MeOH-CH₂Cl₂) gave the title compound (0.055 g, 57%) as a white solid.


M.p.: 148-149°C.

1H-NMR (CDCl₃) δ 7.70 (br s, 1H), 7.27-7.37 (m, 6H),
 20 6.74-6.80 (m, 2H), 3.63 (s, 2H), 3.04 (br d, 2H, J=10.8Hz), 2.88 (t, 2H, J=7.7Hz), 2.05-2.20 (m, 2H), 1.65-1.95 (m, 4H), 1.30-1.60 (m, 3H).

IR (KBr) 3080, 3040, 2945, 1624, 1437, 1384 cm⁻¹.

EIMS: 336.2 (M⁺), 319.2, 255.0, 185.1, 91.1 (base).

25 HRMS calcd. for C₂₁H₂₄N₂O₂: 336.18382. Found: 336.18187.

Example 176-Bromo-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole maleate

30 a) 4-[2-[6-Bromo-1,2-benzisoxazol-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

-66-

The procedure described in Example 1d was followed with 6-bromo-3-methyl-1,2-benzisoxazole (1.02 g, 4.81 mmol), 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (1.72 g, 5.29 mmol), and 1M LDA (5.30 mL, 5.30 mmol) in dry THF (40 mL), except that after addition of reagents, the mixture was stirred at -78°C for 1.5 hours. After purification, the title compound (0.697 g, 35%) was obtained as a pale yellow solid.

¹H-NMR (CDCl₃) δ 7.77 (s, 1H), 7.52 (d, 1H, J=7.2Hz), 7.45 (d, 1H, J=7.2Hz), 4.11 (br d, 2H, J=14.3Hz), 3.02 (t, 2H, J=7.2Hz), 2.70 (dt, 2H, J=12.9Hz, J=3.6Hz), 1.70-1.85 (m, 4H), 1.49 (s, 9H), 1.45-1.55 (m, 1H), 1.09-1.29 (m, 2H).

b) 6-Bromo-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate

The procedure described in Example 1e was followed with the piperidine formed in step a (0.398 g, 0.972 mmol) and TFA (4 mL) in CH₂Cl₂ (26 mL), and triethylamine (1.6 mL, 11.8 mmol) and benzyl bromide (0.155 mL, 1.3 mmol) in CH₂Cl₂ (12 mL). Another reaction with the above piperidine (0.102g, 0.249 mmol) was also carried out. Crude product from both reactions was combined and after purification, the title compound, free base (0.021 g, 4%) was obtained as a tan solid.

The maleate salt was prepared by adding a solution of maleic acid (0.023 g, 0.198 mmol) in EtOH (5 mL) to a solution of the free base (0.072 g, 0.18 mmol) in CH₂Cl₂ (6 mL). After concentration, the residue was recrystallized from EtOH to give the title compound (0.035 g, 38%) as a white solid.

M.p.: 156.8-157.5°C.

¹H-NMR (DMSO-d₆) δ 8.08 (s, 1H), 7.87 (d, 1H, J=8.4Hz), 7.58 (d, 1H, J=8.5Hz), 7.47 (s, 5H), 6.03 (s, 2H), 4.23 (br s, 2H), 3.25-3.40 (m, 2H), 3.02 (t, 2H, J=7.5Hz), 2.80-2.95 (m, 2H), 1.88-2.00 (m, 2H), 1.70-1.80 (m, 2H), 1.30-1.60 (m, 3H).

-67-

^{13}C -NMR (DMSO- d_6) δ 167.2, 162.8, 158.6, 136.0, 131.1, 129.5, 128.9, 127.0, 124.0, 123.7, 120.6, 113.1, 59.6, 51.9, 32.7, 28.9, 21.6.

EIMS: 398 (M⁺), 381, 309, 252, 200, 185 (base), 172,


5 91.

HRMS calc'd for $\text{C}_{21}\text{H}_{23}\text{BrN}_2\text{O}$: 398.0994. Found: 398.0941.

Example 18

6-Cyano-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole

10



15

A solution of NaNO_2 (0.112 g, 1.62 mmol) in H_2O (4 mL) was added to a solution of 6-amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole (0.534 g, 1.59 mmol) in 28% HCl (20 mL) kept at 0°C . The resulting mixture was
20 neutralized to pH 7 by cautious addition of solid Na_2CO_3 . The neutral mixture was added in portions to a well-stirred mixture of toluene (75 mL), ice, and a freshly prepared solution of CuCN (Organic Synthesis, Coll. Vol. I, p. 514; CuSO_4 : 0.318 g, 1.99 mmol). The mixture obtained was kept at
25 0°C for 30 minutes, then at room temperature for 2 hours, and finally, heated at 50°C for 5 minutes. The mixture was extracted with ethyl acetate and the organic phase was washed with H_2O , brine, dried (MgSO_4), filtered, and concentrated. Purification by silica gel flash
30 chromatography (3% $\text{MeOH-CH}_2\text{Cl}_2$) gave the title compound (0.236 g, 43%) as a pale orange solid. Recrystallization (EtOH-hexanes) of a small sample gave the title compound as an off-white solid.

M.p.: $113-114.5^\circ\text{C}$.

35

^1H -NMR (CDCl_3) δ 7.89 (s, 1H), 7.77 (d, 1H, $J=8.4\text{Hz}$), 7.57 (d, 1H, $J=8.7\text{Hz}$), 7.24-7.33 (m, 5H), 3.51 (s, 2H), 3.04

-68-

(t, 2H, J=7.9Hz), 2.92 (br d, 2H, J=11.2Hz), 1.97 (br t, 2H, J=10.7Hz), 1.74-1.85 (m, 4H), 1.26-1.40 (m, 3H).

¹³C-NMR (CDCl₃) δ 161.7, 158.8, 129.2, 128.2, 127.0, 126.3, 125.1, 122.5, 118.1, 114.5, 113.2, 63.4, 53.6, 35.2, 34.0, 32.0, 22.6.


IR (CHCl₃) 2830, 2720, 2160, 1600, 1425, 1385 cm⁻¹.

FABMS: 346 (M⁺ + 1), 309, 275, 239, 155, 119 (base).

HRMS calc'd for C₂₂H₂₃N₃O: 345.1842. Found: 345.1826.

Example 19

10 6-Carboxamide-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole



Powdered KOH (0.150 g, 2.68 mmol) was added to a mixture of 6-cyano-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole (0.250 g, 0.724 mmol) in t-BuOH (10 mL). The resulting mixture was heated at 85°C for 20 minutes. The cooled reaction mixture was poured over brine and extracted with CH₂Cl₂. The organic phase was washed with 10% NaOH, brine, dried (MgSO₄), filtered, and concentrated. The crude product was purified by recrystallization (EtOAc-hexanes) to give the title compound (0.114 g, 43%) as a white solid.


M.p.: 181-182°C.

¹H-NMR (DMSO-d₆) δ 8.20 (br s, 1H, -NH), 8.15 (s, 1H), 7.97 (d, 1H, J=8.4Hz), 7.87 (d, 1H, J=8.1Hz), 7.64 (br s, 1H-NH), 7.22-7.31 (m, 5H), 3.42 (s, 2H), 3.02 (t, 2H, J=7.8Hz), 2.78 (br d, 2H, J=11.5Hz), 1.87 (br t, 2H, J=10.9Hz), 1.69-1.73 (m, 4H), 1.22-1.26 (m, 3H).

¹³C-NMR (DMSO-d₆) δ 167.1, 162.0, 158.9, 138.9, 136.3, 128.7, 126.8, 126.8, 123.3, 122.9, 122.0, 108.7, 62.5, 53.2, 34.9, 33.7, 31.6, 22.0.

FABMS: 346 [(M⁺ + 1), base], 321, 272, 185, 172.

-69-

Example 203-[(1-Phenylmethyl-4-piperidyl)methoxy]-1,2-benzisoxazole
fumarate

a) 4-[(1,2-Benzisoxazol-3-yl)oxymethyl]-1-piperidine-
5 carboxylic acid, 1-(1,1-dimethylethyl)ester

NaH (60% mineral oil dispersion, 0.941 g, 23.53 mmol) was added in portions to a solution of 4-hydroxymethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (4.82 g, 22.41 mmol) in DMF (220 mL) at 0°C. After 10 minutes, the reaction was warmed to room temperature and 3-chloro-1,2-benzisoxazole (3.44 g, 22.41 mmol) was added. The mixture obtained was heated at 115°C for 16 hours. The reaction mixture was diluted with EtOAc and washed with H₂O (4x), brine, dried (MgSO₄), filtered, and concentrated. Purification by silica gel flash chromatography (10% EtOAc-hexanes) gave the title compound (4.16 g, 56%) as a white solid.

M.p.: 103-104.5°C.

¹H-NMR (CDCl₃) δ 7.61 (d, 1H, J=7.9Hz), 7.48-7.54 (m, 1H), 7.41 (d, 1H, J=8.5Hz), 7.22-7.27 (m, 1H), 4.28 (d, 2H, J=6.5Hz), 4.16 (br d, 2H, J=13.3Hz), 2.75 (dt, 2H, J=13.2Hz, J=2.6Hz), 2.04-2.13 (m, 1H), 1.83 (br d, 2H, J=13.7Hz), 1.45 (s, 9H), 1.30 (ddd, 2H, J=24.9Hz, J=12.5Hz, J=4.2Hz).

b) 3-[(1-Phenylmethyl-4-piperidyl)methoxy]-1,2-
25 benzisoxazole fumarate

Trifluoroacetic acid (TFA) (10 mL) was added dropwise to a solution of piperidine formed in step a (0.827 g, 2.49 mmol) in CH₂Cl₂ (25 mL) at 0°C. The resulting mixture was stirred at 0°C for 20 minutes. The mixture was concentrated, and excess TFA was removed by concentrating from toluene. The residue was partitioned between CH₂Cl₂ and

-70-

saturated NaHCO_3 . The organic layer was dried (MgSO_4), filtered, and concentrated. The crude product (0.192 g, 0.827 mmol) was dissolved in CH_2Cl_2 (8 mL), and triethylamine (0.58 mL, 4.13 mmol) followed by benzyl bromide (0.128 mL, 1.07 mmol) was added. The mixture was stirred overnight (18 hours) at room temperature. The reaction was washed with H_2O , brine, dried (MgSO_4), filtered, and concentrated. Purification by silica gel flash chromatography (5% $\text{MeOH}-\text{CH}_2\text{Cl}_2$) gave the title compound, free base (0.199 g, 25%) as an off-white solid.

The fumarate salt was prepared by adding fumaric acid (0.074 g, 0.633 mmol) dissolved in the minimum amount of EtOH to a solution of the free base (0.186 g, 0.58 mmol) in CH_2Cl_2 (6 mL). After concentrating, the crude salt was purified by recrystallization ($\text{EtOH}-\text{Et}_2\text{O}$) to give the title compound (0.134 g, 53%) as an off-white solid.

M.p.: 163-164°C.

$^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ 7.74 (d, 1H, $J=7.8\text{Hz}$), 7.61-7.69 (m, 2H), 7.26-7.41 (m, 6H), 6.60 (s, 2H), 4.28 (d, 2H, $J=6.4\text{Hz}$), 3.65 (s, 2H), 2.97 (br d, 2H, $J=11.4\text{ Hz}$), 2.20 (br t, 2H, $J=11.7\text{Hz}$), 1.90-2.05 (m, 1H), 1.81 (brd, 2H, $J=12.8\text{Hz}$), 1.38-1.49 (m, 2H).

$^{13}\text{C-NMR}$ ($\text{DMSO}-d_6$) δ 166.5, 166.1, 163.3, 137.0, 134.3, 131.1, 129.3, 128.3, 127.3, 123.6, 120.8, 113.5, 110.3, 74.2, 61.7, 52.2, 34.7, 27.6.

IR (KBr) 2980, 2550, 1706, 1650, 1616, 1573, 1447, 1374 cm^{-1} .

EIMS: 305, 185, 172, 91 (base).


HRMS calc'd. for $\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_2$ (freebase): 322.1682. Found: 322.1719.

Anal. Calc'd. for $\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_2 \cdot \text{C}_4\text{H}_4\text{O}_4 \cdot 0.75\text{H}_2\text{O}$: C, 63.78; H, 6.13; N, 6.20. Found: C, 63.88; H, 5.85; N, 6.14.

-71-

Example 213-[(1-Phenylmethyl-4-piperidyl)methylamino]-1,2-benzisoxazole

5



A mixture of 3-chloro-1,2-benzisoxazole (0.238 g, 1.55
10 mmol), 4-aminomethyl-1-phenylmethyl piperidine (0.316 g,
1.55 mmol), and K_2CO_3 (0.214 g, 1.55 mmol) in DMSO (10 mL)
was heated at 150°C for 20 hours. The cooled reaction
mixture was diluted with EtOAc (75 mL) and poured over H_2O
(200 mL). The organic phase was separated and washed with
15 brine, dried ($MgSO_4$), filtered, and concentrated. The brown
oil obtained was purified by silica gel flash chromatography
(4% MeOH- CH_2Cl_2) to give the title compound (0.084 g, 17%) as
pale yellow oil.

1H -NMR ($CDCl_3$): δ 7.44-7.50 (m, 2H), 7.38 (d, 1H,
20 J=8.8Hz), 7.16-7.31 (m, 6H), 4.35-4.42 (m, 1H, -NH-), 3.51
(s, 2H), 3.33 (t, 2H, J=6.2Hz), 2.92 (br d, 2H, J=11.5Hz),
1.99 (t, 2H, J=11.6Hz), 1.77 (br d, 2H, J=12.7Hz), 1.75-1.80
(m, 1H), 1.33-1.45 (m, 2H).

^{13}C -NMR ($CDCl_3$) δ 162.8, 158.6, 137.8, 129.8, 129.3,
25 128.2, 127.2, 122.1, 119.7, 116.2, 110.1, 63.2, 53.3, 49.4,
35.2, 29.9, 29.7.

IR (KBr) 3290, 2924, 2852, 1614, 1564, 1450, 1365 cm^{-1} .


EIMS: 321 (M+), 230, 201, 185, 172, 91 (base).

HRMS calc'd. for $C_{20}H_{23}N_3O$: 321.1842. Found: 321.1825.

-72-

Example 223-[2-[(1-Phenylmethyl)-4-piperidyl]ethylamino]-1,2-benzisoxazole

5



The procedure described in Example 21 was followed with
 10 3-chloro-1,2-benzisoxazole (0.682 g, 4.44 mmol), 4-aminoethyl-1-phenylmethyl piperidine (0.970 g, 4.44 mmol), and K_2CO_3 (0.614 g, 4.44 mmol) in DMSO (30 mL). After purification, the title compound (0.231 g, 15%) was obtained as a pale yellow oil.


15 1H -NMR ($CDCl_3$) δ 7.45-7.54 (m, 2H), 7.24-7.40 (m, 6H), 7.19 (t, 1H, $J=7.8$ Hz), 4.40 (brt, 1H, $J=5.5$ Hz), 3.52 (s, 2H), 3.45 (dt, 2H, $J=7.4$ Hz, $J=6.0$ Hz), 2.91 (brd, 2H, $J=11.8$ Hz), 1.99 (brt, 2H, $J=11.2$ Hz), 1.62-1.73 (m, 4H), 1.37-1.52 (m, 3H).

20 ^{13}C -NMR ($CDCl_3$) δ 162.7, 158.6, 137.8, 129.8, 129.4, 128.2, 127.1, 122.1, 119.8, 116.3, 110.0, 63.3, 53.6, 41.5, 36.1, 33.4, 32.0.

EIMS: 335 (M^+), 244, 199, 186, 172, 91 (base).

HRMS calc'd. for $C_{21}H_{25}N_3O$: 335.1998. Found: 335.1909.

25

Example 233-[3-[1-(Phenylmethyl)-4-piperidyl]propyl]-1,2-benzisoxazole maleate

a) 4-(2-Ethoxy-2-oxoethylidene)-1-piperidine-carboxylic acid, 1-(1,1-dimethylethyl)ester

30 A solution of triethyl phosphonoacetate (3.1 mL, 15.69 mmol) in freshly distilled 1,2-dimethoxyethane (DME, 12.5

-73-

mL) was added to a slurry of NaH (60% mineral oil dispersion, 0.75 g, 18.18 mmol) in DME (6.5 mL). The mixture obtained was stirred for 1 hour at room temperature and a solution of 4-keto-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (2.5 g, 12.55 mmol) in DME (12.5 mL) was added. After stirring overnight (15 hours), the mixture was concentrated. The residue was purified by silica gel flash chromatography (5→20% EtOAc-hexanes) to give the title compound (3.06 g, 91%) as a white solid.

¹H-NMR (CDCl₃) δ 5.71 (s, 1H), 4.16 (q, 2H, J=7.1Hz), 3.45-3.53 (m, 4H), 2.94 (t, 2H, J=5.7Hz), 2.28 (t, 2H, J=5.6Hz), 1.48 (s, 9H), 1.28 (t, 3H, J=7.1Hz).

b) 4-Ethoxycarbonylmethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

A mixture of olefin obtained in step a (3.05 g, 11.3 mmol) and 10% Pd/C (1.2 g, 1.13 mmol) in EtOH (50 mL) was hydrogenated in a Parr shaker at 50 psi for 1.5 hours. The mixture was filtered through a Celite pad, and the filtrate was concentrated to give the title compound (3.07 g, quantitative) as a colorless oil.

¹H-NMR (CDCl₃) δ 4.14 (q, 2H, J=7.0Hz), 4.05-4.12 (m, 2H), 2.72 (br dt, 2H, J=12.1Hz, J=2.8Hz), 2.23 (d, 2H, J=7.2Hz), 1.85-1.95 (m, 1H), 1.61-1.66 (m, 2H), 1.45 (s, 9H), 1.26 (t, 3H, J=7.0Hz), 1.05-1.23 (m, 2H).

c) 4-Hydroxyethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

The procedure described in Example 1b was followed with the ester obtained in step b (3.06 g, 11.3 mmol) and lithium aluminum hydride (0.47 g, 12.4 mmol) in THF (105 mL). Purification by silica gel flash chromatography (50% EtOAc-hexanes) gave unreacted starting material (0.73 g, 24%) and the title compound (1.74 g, 68%) as a colorless oil.

¹H-NMR (CDCl₃) 4.07 (br d, 2H, J=14.1Hz), 3.71 (t, 2H, J=6.5Hz), 2.69 (brt, 2H, J=12.5Hz), 1.50-1.70 (m, 6H), 1.45 (s, 9H), 1.05-1.15 (m, 2H).

-74-

d) 4-Iodoethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

The procedure described in Example 1c was followed with the alcohol obtained in step c (1.74 g, 7.59 mmol),
5 triphenylphosphine (2.49 g, 9.49 mmol), iodine (2.31 g, 9.11 mmol), and pyridine (1.5 mL, 18.2 mmol) in benzene (50 mL). After purification, the title compound (2.27 g, 98%) was obtained as a colorless oil.

¹H-NMR (CDCl₃) δ 4.09 (br d, 2H, J=11.4Hz), 3.22 (t, 2H, J=7.3Hz), 2.70 (brt, 2H, J=12.5Hz), 1.78 (q, 2H, J=6.9Hz),
10 1.47-1.68 (m, 3H), 1.46 (s, 9H), 1.12 (ddd, 2H, J=24.3Hz, J=12.9Hz, J=4.3Hz).

e) 4-[3-[1,2-Benzisoxazol-3-yl]propyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

15 The procedure described in Example 1d was followed with 3-methyl-1,2-benzisoxazole (0.412 g, 3.09 mmol), the iodide obtained in step d (1.15 g, 3.4 mmol), and 1M LDA (3.4 mL, 3.4 mmol) in THF (8 mL). After purification, the title compound (0.694 g, 65%) was obtained as pale yellow oil.

20 ¹H-NMR (CDCl₃) δ 7.65 (d, 1H, J=8.0Hz), 7.54-7.57 (m, 2H), 7.27-7.34 (m, 1H), 4.08 (brd, 2H, J=12.5Hz), 2.99 (t, 2H, J=7.6Hz), 2.66 (brt, 2H, J=12.0Hz), 1.86-1.92 (m, 2H), 1.63-1.68 (m, 4H), 1.45 (s, 9H), 1.36-1.45 (m, 1H), 1.06-1.12 (m, 2H).

25 f) 3-[3-[1-(Phenylmethyl)-4-piperidyl]propyl]-1,2-benzisoxazole maleate

The procedure described in Example 1e was followed with piperidine obtained in step e (0.544 g, 1.58 mmol) and TFA (4 mL) in CH₂Cl₂ (16 mL), and triethylamine (1.1 mL, 7.9
30 mmol) and benzyl bromide (0.21 mL, 1.74 mmol) in CH₂Cl₂ (10 mL) for 7.25 hours. Purification by chromatography (2-5% MeOH-CH₂Cl₂) gave the title compound, free base (0.285 g, 54%) as a pale yellow oil.

The maleate salt was prepared by adding maleic acid
35 (0.109 g, 0.94 mmol) dissolved in the minimum amount of EtOH to a solution of the free base (0.285 g, 0.85 mmol) in CH₂Cl₂ (10 mL). After concentrating, the residue was purified by

-75-

recrystallization (EtOAc) to give the title compound (0.292 g, 76%) as an off-white solid.

M.p.: 134.5-135.5.

¹H-NMR (DMSO-d₆) δ 7.90 (d, 1H, J=7.9Hz), 7.61-7.72 (m, 2H), 7.47 (s, 5H), 7.38 (t, 1H, J=7.3Hz), 6.04 (s, 2H), 4.26 (br s, 2H), 3.22-3.45 (m, 2H), 2.99 (t, 2H, J=7.4Hz), 2.70-2.96 (m, 2H), 1.70-1.90 (m, 4H), 1.40-1.65 (m, 1H), 1.20-1.40 (m, 4H).

¹³C-NMR (DMSO-d₆) δ 167.2, 162.2, 158.4, 135.8, 131.2, 130.3, 129.5, 128.9, 123.5, 122.1, 121.2, 109.7, 59.3, 51.7, 34.8, 32.6, 28.9, 24.5, 24.1.


IR (KBr) 2942, 1705, 1581, 1460, 1359 cm⁻¹.

EIMS: 334 (M⁺, free base), 243, 202, 173 (base), 91.

Anal. calc'd. for C₂₂H₂₆N₂O•C₄H₄O₄: C, 69.31; H, 6.71; N, 6.22. Found: C, 69.11; H, 6.64; N, 6.46.

Example 24

trans-3-[2-[1-(Phenylmethyl)-4-piperidyl]-1,2-benzisoxazole maleate



NaH (60% mineral oil dispersion, 0.10 g, 2.51 mmol) was added to a mixture of 3-triphenylphosphoniummethyl-1,2-benzisoxazole bromide (1.19 g, 2.51 mmol) in THF (10 mL). After stirring at room temperature for 1 hour, a solution of 4-carboxaldehyde-1-phenylmethylpiperidine (0.51 g, 2.51 mmol) in THF (2 mL) was added. The mixture obtained was stirred for 4 hours and filtered. The filtrate was concentrated and the residue was partitioned between diethyl ether and H₂O. The separated organic layer was dried (MgSO₄), filtered, and concentrated. Purification by silica gel flash chromatography (40% EtOAc-hexanes) gave the title compound, free base (0.483 g, 60%) as a colorless oil.

-76-

The maleate salt was prepared by adding maleic acid (0.194 g, 1.67 mmol) dissolved in the minimum amount of EtOH to a solution of the free base (0.483 g, 1.52 mmol) in Et₂O (25 mL). The white solid obtained was collected by
5 filtration to give the title compound (0.581 g, 88%).

M.p.: 174-175°C.

¹H-NMR (DMSO-d₆) δ 8.16 (d, 1H, J=7.8Hz), 7.75 (d, 1H, J=8.4Hz), 7.67 (t, 1H, J=8.2Hz), 7.41-7.55 (m, 6H), 6.96 (br dd, 1H, J=16.5Hz, J=5.6Hz), 6.78 (d, 1H, J=16.5Hz), 6.08 (s,
10 2H), 4.33 (s, 2H), 3.32-3.39 (m, 2H), 2.95-3.15 (m, 2H), 2.50-2.70 (m, 1H), 2.06 (br d, 2H, J=12.6Hz), 1.60-1.90 (m, 2H).

¹³C-NMR (DMSO-d₆) δ 167.3, 162.7, 154.9, 142.7, 135.9, 131.2, 130.4, 130.2, 129.5, 128.9, 124.2, 122.7, 119.4,
15 116.5, 109.9, 59.2, 51.0, 36.2, 27.8.

IR (KBr) 3035, 2944, 1708, 1588, 1472, 1360 cm⁻¹.

EIMS: 3.18 (M⁺, free base), 201, 227, 172, 91 (base).


Anal. Calc'd. for C₂₁H₂₂N₂O⁺·C₄H₄O₄⁻: C, 69.11; H, 6.03; N, 6.45. Found: C, 69.04; H, 6.27; N, 6.38.

20

Example 25

3-[2-[1-(Phenylmethyl)-4-piperazinyl]ethyl]-1,2-benzisoxazole dihydrochloride salt

25



• 2HCl

A mixture of 3-(2-chloroethyl)-1,2-benzisoxazole (0.55 g, 3.03 mmol) and N-benzylpiperazine (1.06 mL, 6.06 mmol) in
30 xylene (4 mL) was heated at 150°C for 4.75 hours. The cooled mixture was diluted with CH₂Cl₂ and washed with H₂O. The organic layer was dried (MgSO₄), filtered and concentrated. Purification by silica gel flash chromatography (50% EtOAc-hexanes-100% EtOAc) gave the title
35 compound, free base (0.337 g, 35%) as a pale yellow oil.

The dihydrochloride salt was made by bubbling excess hydrogen chloride through a solution of the free base (0.337

-77-

g, 1.05 mmol) in Et₂O (50 mL). The white solid formed was collected by filtration to give the title compound (0.259 g, 63%).

M.p.: 233-234°C.

5 ¹H-NMR (D₂O) δ 7.79 (d, 1H, J=8.0Hz), 7.53-7.69 (m, 2H), 7.50 (s, 5H), 7.40 (ddd, 1H, J=7.9Hz, J=6.6Hz, J=1.3Hz), 4.47 (s, 2H), 3.82 (t, 2H, J=7.5Hz), 3.60-3.80 (m, 8H), 3.55 (t, 2H, J=7.5Hz).

¹³C-NMR (D₂O) δ 165.5, 158.0, 134.1, 133.6, 132.4, 130.3,
10 127.1, 124.4, 123.2, 112.8, 63.4, 56.6, 51.8, 51.0, 23.2.

IR (KBr) 2989, 1608, 1436, 1375 cm⁻¹.


EIMS: 321 (M⁺, free base), 256, 189, 91 (base).

Anal. Calc'd. for C₂₀H₂₃N₃O⁺2HCl: C, 60.92; H, 6.39; N, 10.66. Found: C, 60.64; H, 6.57; N, 10.42.

15

Example 26

5,7-Dihydro-7-methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazole-6-one mesylate



NaH (60% mineral oil dispersion, 0.048 g, 1.2 mmol) was
20 added to a solution of 5,7-dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazole-6-one (0.374 g, 1.0 mmol) in DMF (10 mL) at room temperature. After evolution of hydrogen gas had subsided, methyl iodide (0.093 mL, 1.5 mmol) was added and the mixture obtained was
25 stirred for 5.5 hours. Saturated NH₄Cl and H₂O (>50 mL) were added. The reaction was extracted with CH₂Cl₂, and the organic layer was dried (MgSO₄), filtered, and concentrated. Purification by silica gel flash chromatography (CH₂Cl₂-3%

-78-

MeOH-CH₂Cl₂) gave the title compound, free base (0.056 g, 14%) as an off-white foam.

The mesylate salt was made by adding methanesulfonic acid (0.009 mL, 0.144 mmol) to a solution of the free base (0.056 g, 0.044 mmol) in CH₂Cl₂ (5 mL). After concentration, the residue was triturated from Et₂O to give the title compound (0.049 g, 70%) as an off-white solid.

M.p.: 164-165°C (dec).


¹H-NMR (DMSO-d₆) 7.69 (s, 1H), 7.49 (s, 5H), 7.32 (s, 1H), 4.28 (s, 2H), 3.64 (s, 2H), 3.35 (br d, 2H, J=11.6Hz), 3.18 (s, 3H), 2.85-2.99 (m, 4H), 2.30 (s, 3H), 1.95 (br d, 2H, J=12.9Hz), 1.66-1.80 (m, 2H), 1.45-1.60 (m, 1H), 1.35-1.44 (m, 2H).

EIMS: 389 (M⁺, free base), 298, 217, 200, 185 (base), 172.

HRMS Calc'd. for C₂₄H₂₇N₃O₂•CH₃SO₃H: 389.2104. Found: 389.2075.

Example 27

5,7-Dihydro-7-ethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one mesylate



The procedure described in Example 26 was followed with 5,7-dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one (0.374 g, 1.0 mmol), NaH (0.068 g, 1.7 mmol), and ethyl iodide (0.16 mL, 2.0 mmol) in DMF (10 mL). After purification, the title compound, free base (0.076 g, 19%) was obtained as a pale yellow oil.

-79-

The mesylate salt was made by adding methanesulfonic acid (0.007 mL, 0.112 mmol) to a solution of the free base (0.045 g, 0.112 mmol) in CH_2Cl_2 (5 mL). After concentration, the residue was triturated from Et_2O to give the title
 5 compound (0.042 g, 75%) as an off-white solid (hygroscopic).

M.p.: -162°C (dec at $>60^\circ\text{C}$)

$^1\text{H-NMR}$ (DMSO-d_6) δ 7.69 (s, 1H), 7.48 (s, 5H), 7.39 (s, 1H), 4.27 (s, 2H), 3.77 (br q, 2H, $J=7.1\text{Hz}$), 3.64 (s, 2H), 3.34-3.39 (m, 2H), 2.92-2.98 (m, 4H), 2.30 (s, 3H), 1.94 (br
 10 d, 2H, $J=12.8\text{Hz}$), 1.66-1.78 (m, 2H), 1.30-1.60 (m, 3H), 1.16 (t, 3H, $J=7.1\text{Hz}$).

$^{13}\text{C-NMR}$ (DMSO-d_6) 174.5, 163.2, 158.1, 147.0, 131.4, 131.0, 129.8, 129.6, 128.9, 121.2, 117.0, 115.2, 90.0, 59.3, 51.6, 34.5, 34.4, 33.0, 32.6, 28.5, 21.6, 12.3.

15 IR (KBr) 2934, 1716, 1630, 1605, 1466, 1330 cm^{-1} .


EIMS 403 (M^+ , free base), 386, 312, 185 (base), 172.

HRMS Calc'd. for $\text{C}_{25}\text{H}_{29}\text{N}_3\text{O}_2 \cdot \text{CH}_3\text{SO}_3\text{H}$: 403.22605. Found: 403.22761.

Example 28

20

25



5,7-Dihydro-3-[2-[1-(2-chloro-5-thiophenemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one

30 The procedure described in Example 12f was followed with 4-[2-[5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (0.328 g, 0.851 mmol) and TFA (2 mL) in CH_2Cl_2 (8 mL). Only a portion of the crude salt was utilized
 35 in the alkylation step: Na_2CO_3 (0.175 g, 1.66 mmol) and 2-chloro-5-chloromethylthiophene (0.048 mL, 0.40 mmol) in DMF

-80-

(3 mL). After purification (2-4% MeOH-CH₂Cl₂), the title compound (0.0514 g, 43%) was obtained as a white solid.

M.p.: 202-204°C (dec)

¹H-NMR (DMSO-d₆) δ 10.8 (s, 1H), 7.62 (s, 1H), 6.97 (s, 1H), 6.92 (d, 1H, J=3.7Hz), 6.80 (d, 1H, J=3.7Hz), 3.57 (s, 2H), 3.55 (s, 2H), 2.81-2.93 (m, 4H), 1.91 (br t, 2H, J=10.7Hz), 1.61-1.71 (m, 4H), 1.11-1.23 (m, 3H).

¹³C-NMR (DMSO-d₆) δ 176.7, 162.9, 158.3, 146.5, 142.8, 127.0, 126.1, 125.1, 123.2, 117.1, 115.1, 90.1, 57.0, 53.0, 35.0, 34.8, 33.9, 31.6, 21.9.

IR (KBr) 3174, 2950, 1702, 1631, 1453, 1330 cm⁻¹.


EIMS: 398, 382, 350, 322, 236, 172, 91, 81 (base).

HRMS calc'd. for C₂₁H₂₂ClN₃O₂S: 415.1122. Found: 415.1085.

Anal. Calc'd. for C₂₁H₂₂ClN₃O₂S•0.5H₂O: C, 59.36; H, 5.46; N, 9.89. Found: C, 59.21; H, 5.12; N, 9.65.

Example 29

5,7-Dihydro-3-[2-[1-(2-methyl-4-thiazolemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one



The procedure described in Example 12f was followed with 4-[2-[5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (0.367 g, 0.952 mmol) and TFA (2.5 mL) in CH₂Cl₂ (10 mL), and Na₂CO₃ (1.01 g, 9.52 mmol) and 4-chloromethyl-2-methylthiazole hydrochloride salt (0.210 g, 1.142 mmol) in DMF (10 mL). Purification by chromatography (4% MeOH-EtOAc) followed by recrystallization (EtOAc-hexanes) gave the title compound (0.074 g, 20%) as a white solid.

-81-

M.p.: 172-173°C (dec.)

¹H-NMR (CDCl₃) δ 9.40 (br s, 1H), 7.41 (s, 1H), 7.05 (s, 1H), 7.02 (s, 1H), 3.72 (s, 2H), 3.61 (s, 2H), 3.07 (brd, 2H, J=11.3Hz), 2.93 (t, 2H, J=7.8Hz), 2.69 (s, 3H), 2.15 (brt, 2H, J=10.8Hz), 1.76-1.85 (m, 4H), 1.42-1.55 (m, 3H).

¹³C-NMR (CDCl₃) δ 177.4, 166.0, 163.6, 158.3, 151.7, 145.1, 122.2, 116.9, 116.8, 116.3, 91.5, 57.8, 53.4, 35.4, 34.8, 34.0, 31.2, 22.5, 19.2.


IR (KBr) 3101, 3015, 2938, 2924, 1713, 1633, 1456, 1328 cm⁻¹.

EIMS: 396 (M⁺), 379, 284, 267, 206 (base).

HRMS calc'd. for C₂₁H₂₄N₄O₂S: 396.1621. Found: 396.1631.

Example 30

3-[2-[1-(3-Bromophenylmethyl)-4-piperidinyl]ethyl]-5,7-dihydro-6H pyrrolo[4,5-f]-1,2-benzisoxazol-6-one



The procedure described in Example 12f was followed with 4-[2-[5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (0.50 g, 1.30 mmol) and TFA (3 mL) in CH₂Cl₂ (12 mL), and Na₂CO₃ (0.689 g, 6.5 mmol) and 3-bromobenzyl bromide (0.46g, 1.84 mmol) in DMF (20 mL). Purification by chromatography (CH₂Cl₂ → 5% MeOH-CH₂Cl₂) gave the title compound (0.314 g, 53%) as a pale yellow solid. Recrystallization (EtOAc, twice) gave a white solid (0.076 g, 13%).

M.p.: 173-174°C.

¹H-NMR (CDCl₃) δ 8.77 (s, 1H), 7.47 (s, 1H), 7.41 (s, 1H), 7.36 (d, 1H, J=7.7Hz), 7.23 (d, 1H, J=7.9Hz), 7.16 (t,

-82-

1H, J=7.7Hz), 7.00 (s, 1H), 3.61 (s, 2H), 3.46 (s, 2H), 2.86-2.96 (m, 4H), 1.90-2.02 (m, 2H), 1.70-1.85 (m, 4H), 1.30-1.42 (m, 3H).


¹³C-NMR (CDCl₃) δ 177.1, 163.5, 158.4, 144.7, 132.0, 130.1, 129.8, 127.8, 122.3, 122.0, 116.9, 116.5, 91.5, 62.6, 53.6, 35.3, 35.2, 34.3, 31.9, 22.6.

Example 31

3-[2-[1-(4-Bromophenylmethyl)-4-piperidinyl]ethyl]-5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one

10

15



The procedure described in Example 12f was followed with 4-[2-[5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethyl-ethyl)ester (0.50 g, 1.3 mmol) and TFA (3 mL) in CH₂Cl₂ (12 mL), and Na₂CO₃ (0.689 g, 6.5 mmol) and 4-bromobenzyl bromide (0.39 g, 1.56 mmol) in DMF (15 mL). Purification by chromatography (CH₂Cl₂ → 5% MeOH-CH₂Cl₂) gave the title compound (0.415 g, 70%) as an off-white solid.

M.p.: 177-178°C.


¹H-NMR (CDCl₃) δ 9.98 (br s, 1H), 7.37-7.40 (m, 3H), 7.16 (d, 2H, J=8.2Hz), 6.99 (s, 1H), 3.60 (s, 2H), 3.42 (s, 2H), 2.87-2.94 (m, 4H), 1.94 (brt, 2H, J=10.5Hz), 1.65-1.80 (m, 4H), 1.20-1.35 (m, 3H).

¹³C-NMR (CDCl₃) δ 178.1, 163.5, 158.4, 145.2, 137.3, 131.2, 130.8, 122.2, 120.7, 116.7, 116.3, 91.5, 62.5, 53.5, 35.5, 35.2, 34.2, 31.9, 22.5.

-83-

Example 32

5,7-Dihydro-3-[3-[1-(phenylmethyl)-4-piperidinyl]-propyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one



a) 4-[3-[5,7-Dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one-3-yl]propyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

15 The procedure described in Example 7a was followed with 5,7-dihydro-3-methyl-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one (0.13 g, 0.69 mmol), 1M LDA (2.8 mL, 2.8 mmol), and 4-iodoethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (0.233 g, 0.69 mmol) in dry THF (14 mL), except
20 that after addition of reagents, the mixture was stirred at -78°C for 4 hours. Purification by chromatography (10% → 50% EtOAc-CH₂Cl₂) gave recovered starting material (0.031 g, 24%) and the title compound (0.129 g, 47%) as a colorless oil.

25 ¹H-NMR (CDCl₃) δ 9.50 (s, 1H), 7.42 (s, 1H), 7.04 (s, 1H), 4.06 (brd, 2H, J=14.5Hz), 3.62 (s, 2H), 2.91 (t, 2H, J=7.5Hz), 2.66 (dt, 2H, J=13.0Hz, J=2.0Hz), 1.81-1.87 (m, 2H), 1.65 (brd, 2H, J=12.3Hz), 1.44 (s, 9H), 1.34-1.43 (m, 3H), 1.09-1.20 (m, 2H).

30 b) 5,7-Dihydro-3-[3-[3-[1-(phenylmethyl)-4-piperidinyl]propyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one

The procedure described in Example 12f was followed with the piperidine obtained in step a (0.114 g, 0.29 mmol) and TFA (1.5 mL) in CH₂Cl₂ (6 mL), and Na₂CO₃ (0.154 g, 1.45
35 mmol) and benzyl bromide (0.042 mL, 0.35 mmol) in DMF (5 mL). Purification by chromatography (CH₂Cl₂ → 5% MeOH-CH₂Cl₂) gave the title compound (0.070 g, 62%) as a white foamy

-84-

solid. Recrystallization (CH_2Cl_2) gave a white solid (0.054 g, 48%).

M.p.: 164-166°C.

^1H -NMR (CDCl_3) δ 9.73 (br s, 1H), 7.41 (s, 1H), 7.24-7.34 (m, 5H), 6.99 (s, 1H), 3.61 (s, 2H), 3.56 (s, 2H), 2.86-2.97 (m, 4H), 1.97-2.05 (m, 2H), 1.76-1.88 (m, 2H), 1.65-1.70 (m, 2H), 1.26-1.38 (m, 5H).

^{13}C -NMR (CDCl_3) 177.7, 163.6, 158.3, 145.2, 129.5, 128.3, 127.3, 122.2, 116.8, 116.4, 91.6, 63.2, 53.6, 36.1, 35.5, 35.3, 31.9, 25.4, 25.0.


IR (KBr) 3150, 3096, 2930, 1705, 1634, 1495, 1345 cm^{-1} .

EIMS 389 (M^+), 372, 298, 202, 172, 108, 91 (base).

HRMS calc'd. for $\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}$: 389.2104. Found: 389.2107.

Example 33

3-[2-[1-(Phenylmethyl)-4-piperidinylethyl]-5,6,8-trihydro-7H-isoxazolo[4,5-g]quinolin-7-one



a) 6-Acetyl-3,4-dihydro-7-hydroxy-2H-quinolin-2-one

Acetyl chloride (2.0 mL, 28.1 mmol) was added to a mixture of 3,4-dihydro-7-methoxy-2H-quinolin-2-one (1.99 g, 11.2 mmol) in 1,2-dichloroethane (30 mL). The mixture obtained was cooled to 0°C and AlCl_3 (6.0 g, 44.98 mmol) was added in portions. The mixture was heated to reflux for 2 hours. The reaction was cautiously poured over ice- H_2O , stirred for a minimum of 1 hour (to overnight), and extracted with CH_2Cl_2 . The organic layer was washed with brine, dried (MgSO_4), filtered, and concentrated to give the title compound (1.89 g, 82%) as an off-white solid.

-85-

¹H-NMR (DMSO-d₆) δ 12.4 (s, 1H), 10.4 (s, 1H), 7.74 (s, 1H), 6.38 (s, 1H), 2.86 (t, 2H, J=7.4Hz), 2.56 (s, 3H), 2.46-2.56 (m, 2H).

b) 6-Acetyl-3,4-dihydro-7-hydroxy-2H-quinolin-2-one,

5 6-oxime

An aqueous solution of hydroxylamine hydrochloride (1.47 g, 21.2 mmol) and sodium acetate trihydrate (3.0 g, 22.1 mmol) was added to a mixture of the ketone obtained in step a (1.89 g, 9.2 mmol) in EtOH (100 mL). The mixture
10 obtained was heated to reflux for 4 hours. The reaction was concentrated, and the residue was stirred with H₂O. The solid obtained was collected by filtration, and rinsed with EtOH and Et₂O to give the title compound (1.67 g, 82%) as an off-white solid.

15 M.p.: 286.5-287.7°C (dec).

¹H-NMR (DMSO-d₆) δ 11.7 (s, 1H), 11.3 (s, 1H), 10.1 (s, 1H), 7.28 (s, 1H), 6.37 (s, 1H), 2.81 (t, 2H, J=7.5Hz), 2.43 (t, 2H, J=7.5Hz), 2.21 (s, 3H).

c) 6-Acetyl-3,4-dihydro-7-hydroxy-2H-quinolin-2-one,

20 6-oxime acetate

A heterogenous mixture of the oxime obtained in step b (1.67 g, 7.57 mmol) in acetic anhydride (13 mL) was heated at 80°C for 1.5 hours. The reaction mixture was concentrated, and excess acetic anhydride was removed by
25 concentrating from toluene. After drying, the title compound (1.7 g, 86%) was obtained as an off-white solid.

¹H-NMR (DMSO-d₆) δ 11.0 (s, 1H), 10.2 (s, 1H), 7.36 (s, 1H), 6.44 (s, 1H), 2.82 (t, 2H, J=7.5Hz), 2.45 (t, 2H, J=7.5Hz), 2.38 (s, 3H), 2.22 (s, 3H).

30 d) 5,6,8-Trihydro-7H-isoxazolo[4,5-g]quinolin-7-one

A mixture of the oxime acetate obtained in step c (1.58 g, 6.02 mmol) and pyridine (4.9 mL, 60.2 mmol) in DMF (75 mL) was heated in 125-130°C for 2 hours. The reaction was concentrated in vacuo and the residue obtained was purified
35 by recrystallization (EtOAc) to give the title compound (0.80 g, 66%) as a pale yellow solid.

M.p.: 309-311°C (dec).

-86-

¹H-NMR (DMSO-d₆) δ 10.4 (s, 1H), 7.62 (s, 1H), 7.02 (s, 1H), 2.99 (t, 2H, J=7.4Hz), 2.49-2.53 (m, 2H), 2.47 (s, 3H).

e) 4-[2-[5,6,8-Trihydro-7H-isoxazolo[4,5-g]quinolin-7-one-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

The procedure described in Example 7a was followed with the benzisoxazole obtained in step d (0.47 g, 2.3 mmol), 1M LDA (8.1 mL, 8.1 mmol), and 4-iodomethyl-1-piperidine-carboxylic acid, 1-(1,1-dimethylethyl)ester (0.75 g, 2.3 mmol) in dry THF (150 mL), except that after addition of reagents, the mixture was stirred at -78°C for 3.5 hours. An additional reaction with benzisoxazole obtained in step d (0.206 g, 1.02 mmol) was carried out in the same manner. Crude product from both reactions was combined and purification by chromatography (EtOAc) gave the title compound (0.72 g, 54% as a white solid).

¹H-NMR (DMSO-d₆) δ 10.4 (s, 1H), 7.67 (s, 1H), 7.02 (s, 1H), 3.93 (brd, 2H, J=13.4Hz), 2.89-3.31 (m, 4H), 2.57-2.75 (brm, 2H), 2.49-2.53 (m, 2H), 1.64-1.72 (m, 4H), 1.38 (s, 9H), 1.36-1.50 (m, 1H), 1.01-1.16 (m, 2H).

f) 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-5,6,8-trihydro-7H-isoxazolo[4,5-g]quinolin-7-one

The procedure described in Example 12f was followed with the piperidine obtained in step e (0.55 g, 1.37 mmol) and TFA (3.5 mL) in CH₂Cl₂ (14 mL), and Na₂CO₃ (0.758 g, 7.14 mmol) and benzyl bromide (0.23 mL, 1.93 mmol) in DMF (14 mL). After purification by chromatography (5-30% MeOH-CH₂Cl₂), the title compound (0.23 g, 43%) was obtained as a white solid. A sample could be recrystallized from EtOAc-MeOH.

M.p.: 164.4-165.9°C.

¹H-NMR (CDCl₃) δ 9.37 (s, 1H), 7.38 (s, 1H), 7.22-7.36 (m, 5H), 7.01 (s, 1H), 3.55 (s, 2H), 3.09 (t, 2H, J=7.4Hz), 2.91-2.97 (m, 4H), 2.70 (t, 2H, J=7.4Hz), 2.01 (brt, 2H, J=10.3Hz), 1.75-1.80 (m, 4H), 1.39-1.50 (m, 3H).

-87-

^{13}C -NMR (CDCl_3) 170.6, 162.0, 158.2, 141.1, 140.9, 128.7, 128.2, 126.8, 121.2, 120.3, 115.7, 94.6, 62.5, 53.2, 34.9, 33.8, 31.6, 30.2, 24.8, 21.9.


IR (KBr) 3172, 3088, 2921, 1694, 1631, 1447, 1381 cm^{-1} .

5 EIMS 389 (M⁺), 388, 372, 298, 185, 172, 91 (base).

HRMS Calc'd. for $\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_2$: 389.2104. Found: 389.2102.

Example 34

6,8-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-7H-pyrrolo[5,4-g]-1,2-benzisoxazol-7-one



a) 5-Acetyl-1,3-dihydro-4-hydroxy-2H-indol-2-one

An intimate mixture of 4-acetyloxy-1,3-dihydro-2H-indol-2-one (0.876 g, 4.58 mmol) and AlCl_3 (1.83 g, 13.7 mmol) placed in a tear-shape flask was immersed in an oil bath pre-heated to 190°C and heated for 1 hour. Ice-water was added cautiously to the cooled reaction mixture and stirred for 1.5 hours. Concentrated HCl was added and the mixture was extracted with EtOAc . The organic layer was dried (MgSO_4), filtered, and concentrated. Purification by silica gel flash chromatography (1 \rightarrow 3% $\text{MeOH}-\text{CH}_2\text{Cl}_2$) gave the title compound (0.441 g, 50%) as an off-white solid.

^1H -NMR ($\text{DMSO}-d_6$) δ 12.6 (s, 1H), 10.8 (s, 1H), 7.85 (d, 1H, $J=8.4\text{Hz}$), 6.49 (d, 1H, $J=8.4\text{Hz}$), 3.41 (s, 2H), 2.57 (s, 3H).

30 b) 5-Acetyl-1,3-dihydro-4-hydroxy-2H-indol-2-one, 5-oxime

The procedure described in Example 33b was followed with the ketone obtained in step a (0.40 g, 2.09 mmol), and an aqueous solution of NH_2OH hydrochloride (0.29 g, 4.18 mmol), and NaOAc trihydrate (0.569 g, 4.18 mmol) in EtOH (32 mL). After 3 hours, the mixture was diluted with H_2O and extracted with EtOAc . The organic layer was dried (MgSO_4),

-88-

filtered, and concentrated to give the title compound (0.436 g, quantitative) as an off-white solid.

¹H-NMR (DMSO-d₆) δ 12.0 (s, 1H), 11.4 (s, 1H), 10.4 (s, 1H), 7.35 (d, 1H, J=8.2Hz), 6.41 (d, 1H, J=8.3Hz), 3.34 (s, 2H), 2.22 (s, 3H).

c) 5-Acetyl-1,3-dihydro-4-hydroxy-2H-indol-2-one, 5-oxime acetate

The procedure described in Example 33c was followed with the oxime obtained in step b (0.392 g, 1.9 mmol) in Ac₂O (10 mL). The solid obtained was freed from any inorganic salts from previous step by stirring in H₂O. Filtration and drying gave the title compound (0.417 g, 88%) as a red-pinkish solid.

¹H-NMR (DMSO-d₆) δ 11.45 (s, 1H), 10.6 (s, 1H), 7.50 (d, 1H, J=8.2Hz), 6.49 (d, 1H, J=8.2Hz), 3.40 (s, 2H), 2.41 (s, 3H), 2.23 (s, 3H).

d) 6,8-Dihydro-3-methyl-7H-pyrrolo[5,4-g]-1,2-benzisoxazol-7-one

The procedure described in Example 33d was followed with the oxime acetate obtained in step c (0.334 g, 1.35 mmol) and pyridine (0.55 mL, 6.75 mmol) in DMF (25 mL). After work-up, the residue was purified by silica gel flash chromatography (50 → 75% EtOAc-hexanes) to give the title compound (0.086 g, 34%) as a white solid.

M.p.: 259-260°C (dec).

¹H-NMR (DMSO-d₆) δ 10.79 (s, 1H), 7.67 (d, 1H, J=8.5Hz), 6.92 (d, 1H, J=8.3Hz), 3.73 (s, 2H), 2.49 (s, 3H).

e) 4-[2-[6,8-Dihydro-7H-pyrrolo[5,4-g]-1,2-benzisoxazol-7-one-3-yl]ethyl]-1-piperidinecarboxylic acid 1-(1,1-dimethylethyl)ester

The procedure described in Example 7a was followed with the benzisoxazole obtained in step d (0.040 g, 0.213 mmol), 1M LDA (0.85 mL, 0.85 mmol), and 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (0.078 g, 1.234 mmol) in dry THF (20 mL), except that after addition of reagents, the mixture was stirred at -78°C for 4 hours.

-89-

Purification by chromatography (25 → 45% EtOAc-CH₂Cl₂) gave the title compound (0.042 g, 51%) as a pale yellow solid.

¹H-NMR (CDCl₃) δ 8.85 (s, 1H), 7.53 (d, 1H, J=8.1Hz), 6.95 (d, 1H, J=8.3Hz), 4.08-4.14 (m, 2H), 3.78 (s, 2H), 2.99 (t, 2H, J=7.8Hz), 2.68 (brt, 2H, J=12.1Hz), 1.73-1.84 (m, 4H), 1.46-1.60 (m, 1H), 1.46 (s, 9H), 1.17 (ddd, 2H, J=23.2Hz, J=12.1Hz, J=4.3Hz).

f) 6,8-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-7H-pyrrolo[5,4-g]-1,2-benzisoxazol-7-one

The procedure described in Example 12f was followed with the piperidine obtained in step e (0.042 g, 0.109 mmol) and TFA (1.5 mL) in CH₂Cl₂ (6 mL), Na₂CO₃ (0.058 g, 0.545 mmol) and benzyl bromide (0.016 mL, 0.131 mmol) in DMF (6 mL). After purification by chromatography (1-10% MeOH-CH₂Cl₂), the title compound (0.018 g, 44%) was obtained as an off-white solid.

M.p.: 188-189°C.

¹H-NMR (CDCl₃) δ 9.59 (s, 1H), 7.51 (d, 1H, J=8.1Hz), 7.23-7.33 (m, 5H), 6.94 (d, 1H, J=8.2Hz), 3.77 (s, 2H), 3.56 (s, 2H), 2.93-2.99 (m, 4H), 2.02 (brt, 2H, J=10.7Hz), 1.75-1.79 (m, 4H), 1.26-1.39 (m, 3H).


¹³C-NMR (CDCl₃) δ 177.2, 158.9, 158.5, 145.0, 129.4, 128.3, 127.2, 121.5, 117.9, 107.1, 105.1, 63.2, 53.5, 35.1, 34.1, 33.8, 31.7, 29.7, 22.5.

25

Example 35

5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-6H-pyrrolo[5,4-f]-1,2-benzisoxazol-6-one

30



35

-90-

a) 6-Acetyl-1,3-dihydro-5-hydroxy-2H-indol-2-one

The procedure described in Example 33a was followed with 1,3-dihydro-5-methoxy-2H-indol-2-one (4.4 g, 26.96 mmol), acetyl chloride (4.8 mL, 67.41 mmol), and AlCl₃ (14.4 g, 107.8 mmol) in 1,2-dichloroethane (210 mL) for 6 hours. After aqueous work-up and overnight stirring, a precipitate was obtained. This yellow solid was collected by filtration and dried to give the title compound (2.7 g, 52%).

¹H-NMR (DMSO-d₆) δ 12.0 (s, 1H), 10.4 (s, 1H), 7.12 (s, 1H), 6.89 (s, 1H), 3.54 (s, 2H), 2.61 (s, 3H).

b) 6-Acetyl-1,3-dihydro-5-hydroxy-2H-indol-2-one, 6-oxime

The procedure described in Example 33b was followed with the ketone obtained in step a (2.7 g, 14.4 mmol), and an aqueous solution of NH₂OH hydrochloride (2.26 g, 32.5 mmol) and NaOAc trihydrate (4.6 g, 33.9 mmol) in EtOH (155 mL) to give the title compound (2.7 g, 93%) as an off-white solid.

¹H-NMR (DMSO-d₆) δ 11.5 (s, 1H), 11.3 (s, 1H), 10.2 (s, 1H), 6.81 (s, 1H), 6.78 (s, 1H), 3.44 (s, 2H), 2.22 (s, 3H).

c) 6-Acetyl-1,3-dihydro-5-hydroxy-2H-indol-2-one, 6-oxime acetate

The procedure described in Example 33c was followed with the oxime obtained in step b (2.7 g, 13.1 mmol) in Ac₂O (65 mL) for 3 hours to give the title compound (2.93 g, 90%) as an off-white solid.

¹H-NMR (DMSO-d₆) δ 10.4 (s, 1H), 10.2 (s, 1H), 6.86 (s, 1H), 6.81 (s, 1H), 3.48 (s, 2H), 2.37 (s, 3H), 2.22 (s, 3H).

d) 5,7-Dihydro-3-methyl-6H-pyrrolo[5,4-f]-1,2-benzisoxazol-6-one

The procedure described in Example 33d was followed with the oxime acetate obtained in step c (2.75 g, 10.97 mmol) and pyridine (8.9 mL, 109.7 mmol) in DMF (110 mL). After concentration in vacuo, the residue was purified by silica gel flash chromatography (2% MeOH-CH₂Cl₂) to give the title compound (0.245 g, 12%) as a pastel yellow solid.

-91-

¹H-NMR (DMSO-d₆) δ 10.6 (s, 1H), 7.58 (s, 1H), 7.03 (s, 1H), 3.63 (s, 2H), 2.50 (s, 3H).

e) 4-[2-[5,7-Dihydro-6H-pyrrolo[5,4-f]-1,2-benzisoxazol-6-one-3-yl]ethyl]-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester

The procedure described in Example 7a was followed with the benzisoxazole obtained in step d (0.152 g, 0.808 mmol), 1M LDA (3.2 mL, 3.2 mmol), and 4-iodomethyl-1-piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (0.315 g, 0.970 mmol) in dry THF (30 mL), except that after addition of reagents, the mixture was stirred at -78°C for 4.5 hours. Purification by chromatography (50% EtOAc-CH₂Cl₂) gave an inseparable mixture (0.153 g, 1.6:1) of starting material and the title compound, respectively, as a pale yellow soft solid. In a separate experiment, a better ratio (starting material/title compound → 1:3) was observed.

¹H-NMR (CDCl₃) δ 9.84 (s, 1H), 7.45 (s, 1H), 7.05 (s, 1H), 4.05-4.15 (m, 2H), 3.69 (s, 2H), 2.96 (t, 2H, J=7.8Hz), 2.68 (brt, 2H, J=11.8Hz), 1.72-1.82 (m, 4H), 1.45 (s, 9H), 1.43-1.53 (m, 1H), 1.15 (ddd, 2H, J=23.6Hz, J=12.1Hz, J=4.0Hz).

f) 5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-6H-pyrrolo[5,4-f]-1,2-benzisoxazol-6-one

The procedure described in Example 12f was followed with the mixture obtained in step a (0.153 g) and TFA (1.5 mL) in CH₂Cl₂ (6 mL). Only a portion of the crude salt mixture (0.081 g) was utilized in the alkylation step: Na₂CO₃ (0.041 g, 0.388 mmol) and benzyl bromide (0.012 mL, 0.101 mmol) in DMF (6 mL). Note: For the alkylation step, the solvent (DMF) was thoroughly degassed with Ar and work-up with saturated NaHCO₃ was omitted (brine was used instead). After purification (2-6% MeOH-CH₂Cl₂), the title compound (0.018 g, 60% based on percentage of desired starting material) was obtained as a light yellow solid.

M.p.: 204.5-205.5°C (dec).


¹H-NMR (CDCl₃) δ 8.09 (s, 1H), 7.46 (s, 1H), 7.27-7.33 (m, 5H), 6.99 (s, 1H), 3.68 (s, 2H), 3.50-3.52 (m, 2H),

-92-

2.90-2.99 (m, 4H), 1.92-2.05 (m, 2H), 1.70-1.80 (m, 4H),
1.30-1.40 (m, 3H).

Example 36

5 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1H-indazole
maleate



a) 1-(2-Fluorophenyl)-3-[1-(phenylmethyl)-4-
piperidinyl]propanone

Lithium hexamethyldisilazide (1M, 14.5 mL, 14.5 mmol)
was added to a solution of o-fluoroacetophenone (2.0 g, 14.5
10 mmol) in THF (100 mL) at -78°C. The mixture obtained was
allowed to warm to -20°C over a 30-60 minute period and then
re-cooled to -78°C. A solution of 4-carboxaldehyde-1-
piperidinecarboxylic acid, 1-(1,1-dimethylethyl)ester (2.94
g, 14.5 mmol) in THF (20 mL) was added. The mixture was
15 kept at -78°C for 30 minutes and then allowed to warm to
room temperature and stirred for 40 minutes. Saturated NH₄Cl
was added and the reaction was extracted with EtOAc. The
organic layer was dried (MgSO₄), filtered, and concentrated.
Purification by silica gel flash chromatography (30 → 50%
20 EtOAc-CH₂Cl₂) gave a yellow oil (2.14 g). ¹H-NMR showed
mixture of compounds (two major components). Further
purification was not attempted. Used as such in next step.

A portion of the mixture obtained above (0.661 g) and
PtO₂ (0.070 g, 0.31 mmol) in EtOH was hydrogenated in a Parr
25 shaker at 48 psi for 2 hours. The mixture was filtered
through a Celite pad and the filtrate was concentrated. An
additional reaction with the mixture above (1.00 g) was also
carried out. Crude products from these reactions were
combined and purified by silica gel flash chromatography (1
30 → 3% MeOH-CH₂Cl₂, then 30% EtOAc-hexanes) to give the title
compound (0.192 g, 5.3% overall) as a colorless oil.

-93-

¹H-NMR (CDCl₃) δ 7.85 (dt, 1H, J=7.6Hz, J=1.8Hz), 7.47-7.56 (m, 1H), 7.09-7.33 (m, 7H), 3.49 (s, 2H), 2.95-3.03 (m, 2H), 2.89 (brd, 2H, J=11.4Hz), 1.94 (brt, 2H, J=11.1Hz), 1.63-1.77 (m, 4H), 1.24-1.39 (m, 3H).

5 EIMS: 325 (M⁺), 202, 188, 172, 91, 66 (base).

b) 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1H-indazole maleate

A mixture of the ketone obtained in step b (0.178 g, 0.55 mmol) in anhydrous hydrazine (10 mL) was heated to
10 reflux for 3 hours. The reaction was allowed to cool, H₂O was added, and the mixture was extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), filtered, and concentrated. Purification by silica gel radial chromatography (CH₂Cl₂ →
10% MeOH-CH₂Cl₂) gave the title compound, free base (0.047 g,
15 27%) as a colorless oil.

The maleate salt was prepared by adding a solution of maleic acid (0.010 g, 0.085 mmol) in Et₂O (5 mL) to a solution of the free base (0.027 g, 0.085 mmol) in Et₂O (75 mL). The white solid obtained was collected by filtration
20 to give the title compound (0.014 g, 38%).


M.p.: 151.5-152.5°C

¹H-NMR (DMSO-d₆) δ 12.64 (s, 1H), 7.72 (d, 1H, J=8.0Hz), 7.44-7.47 (m, 6H), 7.31 (t, 1H, J=7.3Hz), 7.06 (t, 1H, J=7.4Hz), 6.02 (s, 2H), 4.24 (br s, 2H), 2.93 (t, 2H, J=7.6Hz), 1.90-2.00 (m, 2H), 1.60-1.80 (m, 3H), 1.30-1.50 (m, 4H).

-94-

CLAIMS

1. A compound of the formula




- 10 wherein R^1 and R^2 are independently selected from hydrogen, (C_1-C_6) alkoxy, benzyloxy, phenoxy, hydroxy, phenyl, benzyl, halo, nitro, cyano, COR^5 , $-COOR^5$, $-CONHR^5$, $-NR^5R^6$, $-NR^5COR^6$, $-OCONR^5R^6$, $-NHCOOR^5$, (C_1-C_6) alkyl optionally substituted with from 1 to 3 fluorine atoms; SO_pCH_2 -phenyl or $SO_p(C_1-C_6)$ alkyl,
- 15 wherein p is 0, 1 or 2; pyridylmethyloxy or thienylmethyloxy; 2-oxazolyl, 2-thiazolyl and benzenesulfonamide; wherein the phenyl moieties of said phenoxy, benzyloxy, phenyl, benzyl and benzenesulfonamide groups, the pyridyl and thienyl moieties of said pyridylmethyloxy or
- 20 thienylmethyloxy and the oxazolyl and thiazolyl moieties of said 2-oxazolyl and 2-thiazolyl may optionally be substituted with 1 or 2 substituents independently selected from halo, (C_1-C_4) alkyl, trifluoromethyl, (C_1-C_4) alkoxy, cyano, nitro and hydroxy;
- 25 or R^1 and R^2 , when attached to adjacent carbon atoms and when X is oxygen, sulfur or NR^4 wherein R^4 is hydrogen or (C_1-C_4) alkyl) may form, together with the carbon atoms to which they are attached, a group of the formula

30

35

-95-



10 wherein J is oxygen, sulfur or NR^4 , "a" is 1 or 2, R^3 is hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$ and Q is oxygen, sulfur, NH , CHCH_3 , $\text{C}(\text{CH}_3)_2$, $-\text{CH}=\text{CH}-$, or $(\text{CH}_2)_1$ wherein 1 is an integer from 1 to 3;

X is oxygen, sulfur, $-\text{CH}=\text{CH}-$, $-\text{CH}=\text{N}-$, $-\text{N}=\text{CH}-$, $-\text{N}=\text{N}-$, or
 15 NR^4 wherein R^4 is hydrogen or $(\text{C}_1\text{-C}_4)\text{ alkyl}$;

Y is $-(\text{CH}_2)_m-$, $-\text{CH}=\text{CH}(\text{CH}_2)_n-$, $-\text{NR}^4(\text{CH}_2)_m-$, or $-\text{O}(\text{CH}_2)_m-$ wherein R^4 is defined as above, n is an integer from 0 to 3 and m is an integer from 1 to 3;


R^5 and R^6 are each independently selected from hydrogen,
 20 $(\text{C}_1\text{-C}_6)\text{alkyl}$, phenyl or benzyl, wherein the phenyl moieties of said phenyl and benzyl may optionally be substituted with 1 or 2 substituents independently selected from fluoro, chloro, bromo, iodo, $(\text{C}_1\text{-C}_4)\text{ alkyl}$, trifluoromethyl, $(\text{C}_1\text{-C}_4)\text{ alkoxy}$, cyano, nitro and hydroxy, or NR^5R^6 together form a 4
 25 to 8 membered ring wherein one atom of the ring is nitrogen and the others are carbon, oxygen or nitrogen, or NR^5COR^6 together form a 4 to 8 membered cyclic lactam ring;

M is $-\text{CH}-$ or nitrogen;

L is phenyl, phenyl- $(\text{C}_1\text{-C}_6)\text{alkyl}$, cinnamyl or
 30 pyridylmethyl, wherein the phenyl moieties of said phenyl and phenyl- $(\text{C}_1\text{-C}_6)\text{alkyl}$ may optionally be substituted with 1-3 substituents independently selected from $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_1\text{-C}_4)\text{alkoxycarbonyl}$, $(\text{C}_1\text{-C}_4)\text{alkylcarbonyl}$ - OCONR^5R^6 , $-\text{NHCOOR}^5$ or halo; or L is a group of the formula

35

-96-



5

wherein b is an integer from 1 to 4, R^{13} and R^{14} are independently selected from hydrogen, (C_1-C_4) alkyl, halo and phenyl, E and F are independently selected from -CH- and nitrogen, and G is oxygen, sulfur or NR^4 wherein R^4 is defined as above, with the proviso that when E and F are both nitrogen, one of R^{13} and R^{14} is absent; and

R^7 and R^8 are independently selected from hydrogen, (C_1-C_6) alkyl, (C_1-C_6) alkoxycarbonyl, (C_1-C_6) alkylcarbonyl and (C_1-C_6) alkoxy, with the proviso that said (C_1-C_6) alkoxy is not attached to a carbon that is adjacent to a nitrogen;

or pharmaceutically acceptable salt of such compound.

2. A compound according to claim 1, wherein X is sulfur or oxygen, Y is $-CH_2-CH_2-$, or $-CH_2-$, M is -CH- and L is benzyl, R^1 and R^2 are (C_1-C_6) alkyl, (C_1-C_6) alkoxy, NR^5R^6 , or NR^5COR^6 , R^3 is hydrogen or (C_1-C_6) alkyl, J is oxygen or sulfur and Q is $CH(CH_3)$, $CH(CH_3)_2$, $-CH=CH$ or $(CH_2)_1$.

3. A compound selected from the group consisting of:

3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5-Methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5,6-Dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

7-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-Acetamido-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-Amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-Benzamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5 6-Benzenesulfonamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

6-(4-Morpholinyl)-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;

10 1-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-isoquinoline;

3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisothiazole;

15 4-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,3-quinazoline;

6-Hydroxy-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;

20 6-Bromo-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;

6-Cyano-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;

6-Carboxamide-3-[2-[1-(phenylmethyl)-4-piperidyl]ethyl]-1,2-benzisoxazole;

25 3-[(1-phenylmethyl-4-piperidyl)methoxy]-1,2-benzisoxazole;

3-[(1-phenylmethyl-4-piperidyl)methylamino]-1,2-benzisoxazole;

30 3-[2-(1-phenylmethyl)-4-piperidyl]ethylamino]-1,2-benzisoxazole;

3-[3-[1-(phenylmethyl)-4-piperidyl]propyl]-1,2-benzisoxazole;

trans-3-[2-[1-(phenylmethyl)-4-piperidyl]ethenyl]-1,2-benzisoxazole;

35 3-[2-[1-(phenylmethyl)-4-piperazinyl]ethyl]-1,2-benzisoxazole;

-98-

- 5,7-Dihydro-7-methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
5,7-Dihydro-7-ethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
5 5,7-Dihydro-3-[2-[1-(2-chloro-5-thiophenemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
5,7-Dihydro-3-[2-[1-(2-methyl-4-thiazolemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
3-[2-[1-(3-Bromophenylmethyl)-4-piperidinyl]-5,7-
10 dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
3-[2-[1-(4-Bromophenylmethyl)-4-piperidinyl]ethyl]-5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
5,7-Dihydro-3-[3-[1-(phenylmethyl)-4-piperidinyl]-propyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
15 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-5,6,8-trihydro-7H-isoxazolo[4,5-g]quinolin-7-one;
6,8-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-7H-pyrrolo[5,4-g]-1,2-benzisoxazol-7-one;
5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-6H-pyrrolo[5,4-f]-1,2-benzisoxazol-6-one;
20 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1H-indazole;

and the pharmaceutically acceptable salts of such compounds.

- 25 4. A compound according to claim 1, selected from the group consisting of:


- 3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate;
5,6-dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole maleate;
30 5-methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate;
7-methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole fumarate;
35 6-benzamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole maleate;

-99-

6-benzenesulfonamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole fumarate; and

1-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]isoquinoline maleate.


5 5. A compound of the formula



10

wherein W is a leaving group; j is an integer from 0 to 2; R¹⁰ is a nitrogen protecting group; and R⁷ and R⁸ are independently selected from hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkylcarbonyl and (C₁-C₆)alkoxy, with the proviso that said (C₁-C₆)alkoxy is not attached to a carbon that is adjacent to a nitrogen.

6. A compound of the formula




20

25

wherein R¹, R², R⁷, R⁸, X, Y and M are as defined in claim 1 and R¹¹ is hydrogen or a nitrogen protecting group.

7. A compound of the formula



30

35

-100-

wherein R^1 , R^2 , R^7 , R^8 and L are as defined in claim 1 and Y' is $-\text{CH}=\text{CH}-(\text{CH}_2)_n-$ or $-(\text{CH}_2)_m-$.


8. A pharmaceutical composition for enhancing memory or treating or preventing Alzheimer's disease comprising a compound according to claim 1 and a pharmaceutically acceptable carrier.

9. A pharmaceutical composition for inhibiting cholinesterase in a mammal, comprising a compound according to claim 1 and a pharmaceutically acceptable carrier.

10. A method of inhibiting cholinesterase in a mammal, comprising administering to a mammal a cholinesterase inhibiting amount of a compound according to claim 1.

11. A method of enhancing memory or treating or preventing Alzheimer's disease, comprising administering to a patient a memory enhancing effective amount of a compound according to claim 1.

12. A process for preparing a compound having the formula



I-A


wherein R^1 and R^2 are independently selected from hydrogen, (C_1-C_6) alkoxy, benzyloxy, phenoxy, hydroxy, phenyl, benzyl, halo, nitro, cyano, COR^5 , $-\text{COOR}^5$, $-\text{CONHR}^5$, $-\text{NR}^5\text{R}^6$, $-\text{NR}^5\text{COR}^6$, $-\text{OCONR}^5\text{R}^6$, $-\text{NHCOOR}^5$, (C_1-C_6) alkyl optionally substituted with from 1 to 3 fluorine atoms; SO_pCH_2 -phenyl or $\text{SO}_p(\text{C}_1-\text{C}_6)$ alkyl, wherein p is 0, 1 or 2; pyridylmethyloxy or thienyl-methyloxy; 2-oxazolyl, 2-thiazolyl and benzenesulfonamide; wherein the phenyl moieties of said phenoxy, benzyloxy, phenyl, benzyl and benzenesulfonamide groups, the pyridyl and thienyl moieties of said pyridylmethyloxy or thienylmethyloxy and the oxazolyl and

-101-

thiazolyl moieties of said 2-oxazolyl and 2-thiazolyl may optionally be substituted with 1 or 2 substituents independently selected from halo, (C₁-C₄)alkyl, trifluoromethyl, (C₁-C₄)alkoxy, cyano, nitro and hydroxy;

5 or R¹ and R², when attached to adjacent carbon atoms and when X is oxygen, sulfur or NR⁴ wherein R⁴ is hydrogen or (C₁-C₄ alkyl) may form, together with the carbon atoms to which they are attached, a group of the formula

10



15

wherein J is oxygen, sulfur or NR⁴, "a" is 1 or 2, R³ is hydrogen or (C₁-C₆)alkyl and Q is oxygen, sulfur, NH, CHCH₃,
 20 C(CH₃)₂, -CH=CH-, or (CH₂)₁ wherein 1 is an integer from 1 to 3;

X is oxygen, sulfur, -CH=CH-, -CH=N-, -N=CH-, -N=N-, or NR⁴ wherein R⁴ is hydrogen or (C₁-C₄) alkyl;

m is an integer from 1 to 3;


25 R⁷ and R⁸ are independently selected from hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkylcarbonyl and (C₁-C₆)alkoxy, with the proviso that said (C₁-C₆)alkoxy is not attached to a carbon that is adjacent to a nitrogen;

L is phenyl-(C₁-C₆)alkyl, pyridylmethyl, or a group of
 30 the formula K;

or a pharmaceutically acceptable acid addition salt thereof; comprising: (a) reacting a compound of formula

35


-102-



VI

10 wherein R^1 , R^2 , X , m , R^7 , and R^8 are defined as above
 with a compound of formula WL wherein W is a leaving group
 and L is defined as above; and (b) optionally reacting the
 compound of formula I-A obtained thereby with a
 pharmaceutically acceptable salt.


15 13. A process for preparing a compound having the
 formula



I-B

25 wherein R^1 , R^2 , X , m , R^7 and R^8 are defined as in claim 12;
 L is phenyl or cinnamyl;
 or a pharmaceutically acceptable acid addition salt
 thereof;

comprising: (a) reacting a compound of formula



II


-103-

wherein R^1 , R^2 , and X are defined as in claim 12 with a compound of the formula



10 wherein W , m , R^7 and R^8 are defined as in claim 12, and R^{10} is L as defined above; and (b) optionally reacting the compound of formula I-B obtained thereby with a pharmaceutically acceptable salt.

14. A process for preparing a compound having the
15 formula



wherein R^1 , R^2 , X , R^7 and R^8 are defined as in claim 12;

n is an integer from 0 to 3;

25 M is nitrogen or $-CH-$;

L is phenyl, cinnamyl, phenyl- (C_1-C_6) alkyl, pyridylmethyl, or a group of the formula K;

or a pharmaceutically acceptable acid addition salt thereof;


30 comprising: (a) reacting a compound of formula



-104-

wherein R^1 , R^2 and X are defined as in claim 12, R^9 is hydrogen or halo with a compound of the formula


5



10 wherein R^7 and R^8 are defined as in claim 12, and n , M and L are defined as above; and (b) optionally reacting the compound of formula I-C obtained thereby with a pharmaceutically acceptable salt.

15 15. A process for preparing a compound having the formula

20



wherein R^1 , R^2 , X , R^7 , R^8 and L are defined as in claim 12;

25 M is defined as in claim 14;


Y is $-(CH_2)_m-$, $-CH=CH(CH_2)_n-$, $-NR^4(CH_2)_m-$, or $-O(CH_2)_m-$ wherein R^4 is defined as above, n is an integer from 0 to 3 and m is an integer from 1 to 3;

or a pharmaceutically acceptable acid addition salt
30 thereof;

comprising: (a) reacting a compound of formula


35

-105-



XI


wherein R^1 , R^2 and X are defined as in claim 12, R^{12} is halo
 10 with a compound of the formula



XII

wherein m , R^7 and R^8 are defined as in claim 12, M and L are
 20 defined as in claim 14, and Z is OH or NHR^4 wherein R^4 is
 hydrogen or (C_1-C_4) alkyl; and (b) optionally reacting the
 compound of formula I-D obtained thereby with a
 pharmaceutically acceptable salt.

16. A process for preparing a compound having the
 25 formula




I-D

wherein R^1 , R^2 , X , R^7 and R^8 are defined as in claim 12;
 35 M and L are defined as in claim 14;
 Y is defined as in claim 15;

-106-


or a pharmaceutically acceptable acid addition salt thereof;

comprising: (a) reacting a compound of formula



wherein R^1 , R^2 , X , R^7 and R^8 are defined as in claim 12, M is defined as in claim 3, and Y is defined as in claim 15 with a compound of the formula WL wherein W and L are defined as in claim 12, and (b) optionally reacting the compound of formula I-D obtained thereby with a pharmaceutically acceptable salt.

17. A process for preparing a compound having the formula



wherein R^1 , R^2 , X , m , R^7 and R^8 are defined as in claim 12;


L is defined as in claim 14;

30 or a pharmaceutically acceptable acid addition salt thereof;

comprising: (a) reacting a compound of formula


35

-107-



XIII


wherein R^1 , R^2 and X are defined as in claim 12 and R^{12} is halo with a compound of the formula



XIV

wherein R^7 and R^8 are defined as in claim 12, and L is defined as in claim 14; and (b) optionally reacting the compound of formula I-E obtained thereby with a pharmaceutically acceptable salt.

18. A process for preparing a compound having the formula



I-F

wherein R^1 , R^2 , R^7 and R^8 are defined as in claim 12;

L is defined as in claim 14;

X is oxygen or NR^4 wherein R^4 is defined as in claim 15;


n' is an integer from 0 to 3;

or a pharmaceutically acceptable acid addition salt thereof;

comprising: (a) reacting a compound of formula

-108-


5



wherein R^1 , R^2 , R^7 and R^8 are defined as in claim 12, L is
 10 defined as in claim 14, and n' is defined as above with
 hydrazines or hydroxylamines; and (b) optionally reacting
 the compound of formula I-F obtained thereby with a
 pharmaceutically acceptable salt.

19. A process for preparing a compound having the
 15 formula

20




25 wherein R^1 , R^2 , R^7 and R^8 are defined as in claim 12;
 L is defined as in claim 14;
 X is defined as in claim 18;
 n' is an integer from 0 to 1;
 or a pharmaceutically acceptable acid addition salt
 30 thereof;
 comprising: (a) reacting a compound of formula

35

-109-


5



10 wherein R^1 , R^2 , R^7 and R^8 are defined as in claim 12, L is defined as in claim 14, and n' is defined as above with hydrazines or hydroxylamines; and (b) optionally reacting the compound of formula I-G obtained thereby with a pharmaceutically acceptable salt.

15 20. A process for preparing a compound having the formula

20



I-H

25


wherein R^1 , R^2 , m , R^7 , R^8 and L are defined as in claim 12;
or a pharmaceutically acceptable acid addition salt thereof;

comprising: (a) reacting a compound of formula

30

35


-110-



VI'

10 wherein R¹, R², m, R⁷ and R⁸ are defined as in claim 12 with
a compound of formula WL wherein W and L are defined as in
claim 12; and (b) optionally reacting the compound of
formula I-H obtained thereby with a pharmaceutically
acceptable salt.

15 21. A process for preparing a compound having the
formula



25

wherein m, R⁷, R⁸ and L are defined as in claim 12;

X and J are oxygen, sulfur, or NR⁴ wherein R⁴ is defined
as in claim 15;


Q is oxygen, sulfur, NH, CHCH₃, C(CH₃)₂, -CH=CH-, or
30 (CH₂)₁ wherein 1 is an integer from 1 to 3;

or a pharmaceutically acceptable acid addition salt
thereof;

comprising: (a) reacting a compound of formula


35

-111-



10 wherein m , R^7 and R^8 are defined as in claim 12, and X , J and q are defined as above with a compound of formula WL wherein W and L are defined as in claim 12; and (b) optionally reacting the compound of formula I-J obtained thereby with a pharmaceutically acceptable salt.

15 22. A process for preparing a compound having the formula



25

wherein m , R^7 and R^8 are defined as in claim 12;

L is defined as in claim 14;

X , J and Q are defined as in claim 21;


30 or a pharmaceutically acceptable acid addition salt thereof;

comprising: (a) reacting a compound of formula

35

-112-


5



I-A'

10 wherein m, R⁷, R⁸ and L are defined as in claim 12, X, J, and Q are defined as in claim 21, or a compound of the formula

15




I-B'

20

wherein m, R⁷, and R⁸ are define as in claim 12, L is defined as in claim 13, X, J and Q are defined as in claim 21 with an alkylating agent of formula (C₁-C₄)alkyl-W wherein W is defined as in claim 12; and (b) optionally reacting the
25 compound of formula I-J obtained thereby with a pharmaceutically acceptable salt.

23. A process for preparing a compound having the formula

30



I-I

35

-113-


wherein m, R⁷, R⁸ and L are defined as in claim 12;

X and Q are defined as in claim 21;

R³ is (C₁-C₆)alkyl;

or a pharmaceutically acceptable acid addition salt thereof;

comprising: (a) reacting a compound of formula



wherein m, R⁷ and R⁸ are defined as in claim 12, X and Q are defined as in claim 21, and R³ is defined as above with a compound of the formula WL wherein W and L are defined as in claim 12; and (b) optionally reacting the compound of formula I-I obtained thereby with a pharmaceutically acceptable salt.

24. A process according to any of the preceding claims wherein X is sulfur or oxygen, Y is -CH₂-CH₂-, or -CH₂-, M is -CH- and L is benzyl, R¹ and R² are (C₁-C₆)alkyl, (C₁-C₆)alkoxy, NR⁵R⁶, or NR⁵COR⁶, R³ is hydrogen or (C₁-C₆)alkyl, J is oxygen or sulfur and Q is CH(CH₃), CH(CH₃)₂, -CH=CH or (CH₂)₁.

25. A process according to any of the preceding claims wherein said compound is selected from the group consisting of:

3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5-Methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

5,6-Dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;

-114-

- 5-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 6-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 5 7-Methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 6-Acetamido-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 6-Amino-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 10 1,2-benzisoxazole;
- 6-Benzamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 6-Benzenesulfonamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 15 6-(4-Morpholinyl)-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 1-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-isoquinoline;
- 20 3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisothiazole;
- 4-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,3-quinazoline;
- 25 6-Hydroxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 6-Bromo-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 6-Cyano-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 30 6-Carboxamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole;
- 3-[1-(phenylmethyl)-4-piperidinyl]methoxy]-1,2-benzisoxazole;
- 35 3-[1-(phenylmethyl)-4-piperidinyl]methylamino]-1,2-benzisoxazole;

-115-

- 3-[2-(1-Phenylmethyl)-4-piperidyl]ethylamino]-1,2-benzisoxazole;
- 3-[3-[1-(Phenylmethyl)-4-piperidyl]propyl]-1,2-benzisoxazole;
- 5 trans-3-[2-[1-(Phenylmethyl)-4-piperidyl]ethenyl]-1,2-benzisoxazole;
- 3-[2-[1-(Phenylmethyl)-4-piperazinyl]ethyl]-1,2-benzisoxazole;
- 5,7-Dihydro-7-methyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 10 5,7-Dihydro-7-ethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 5,7-Dihydro-3-[2-[1-(2-chloro-5-thiophenemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 15 5,7-Dihydro-3-[2-[1-(2-methyl-4-thiazolemethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 3-[2-[1-(3-Bromophenylmethyl)-4-piperidinyl]-5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 3-[2-[1-(4-Bromophenylmethyl)-4-piperidinyl]ethyl]-5,7-dihydro-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 20 5,7-Dihydro-3-[3-[1-(phenylmethyl)-4-piperidinyl]propyl]-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one;
- 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-5,6,8-trihydro-7H-isoxazolo[4,5-g]quinolin-7-one;
- 25 6,8-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-7H-pyrrolo[5,4-g]-1,2-benzisoxazol-7-one;
- 5,7-Dihydro-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-6H-pyrrolo[5,4-f]-1,2-benzisoxazol-6-one;
- 3-[2-[1-(Phenylmethyl)-4-piperidinyl]ethyl]-1H-indazole;
- 30 and the pharmaceutically acceptable salts of such compounds.
26. A process according to any of the preceding claims wherein said compound is selected from the group consisting of:
- 35 3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate;

-116-

5,6-dimethyl-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole maleate;

5-methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole maleate;

5 7-methoxy-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole fumarate;

6-benzamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]-ethyl]-1,2-benzisoxazole maleate;


6-benzenesulfonamide-3-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]-1,2-benzisoxazole fumarate; and

10 1-[2-[1-(phenylmethyl)-4-piperidinyl]ethyl]isoquinoline maleate.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 92/01605

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all)		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.C1.5	C 07 D 413/06	A 61 K 31/445
C 07 D 401/06	C 07 D 417/06	C 07 D 261/20
C 07 D 211/32		C 07 D 413/12
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.C1.5	C 07 D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸ /		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	EP,A,0296560 (EISAI CO., LTD) 28 December 1988, see the claims ---	1,8-11
Y	EP,A,0091511 (HOECHST ROUSSEL) 19 October 1983, see the claims ---	1,8-11
Y	EP,A,0229391 (EISAI CO., LTD) 22 July 1982, see the claims ---	1,8-11
P,X	EP,A,0428437 (ADIR ET COMPAGNIE) 22 May 1991, see the claims -----	1,8-11
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
23-07-1992	19.08.92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE		

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Remark: Although claims 10 and 11 are directed to a method of treatment of human body, the search has been carried out and based on the alleged effects of the compounds.

2. ☒ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Remark: In claim 5 the definition of the variables W and R₁₀ are vague and do not concisely define the matter for which protection is sought so that a comprehensive search is not possible

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 9201605

SA 60166

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 12/08/92. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0296560	28-12-88	JP-A- 1079151	24-03-89
		US-A- 4895841	23-01-90
		US-A- 5100901	31-03-92
EP-A- 0091511	19-10-83	US-A- 4390544	28-06-83
		US-A- 4536578	20-08-85
		AU-B- 560453	09-04-87
		AU-A- 8977882	13-10-83
		CA-A- 1209146	05-08-86
		DE-A- 3278471	16-06-88
		JP-A- 58177980	18-10-83
		US-A- 4604395	05-08-86
		US-A- 4610988	09-09-86
EP-A- 0229391	22-07-87	US-A- 4609658	02-09-86
		AU-A- 6690686	02-07-87
		CA-A- 1279317	22-01-91
		JP-A- 62234065	14-10-87
		US-A- 4942169	17-07-90
		US-A- 5039681	13-08-91
		US-A- 5118684	02-06-92
EP-A- 0428437	22-05-91	US-A- 4849431	18-07-89
		FR-A- 2654104	10-05-91
		AU-A- 6581990	16-05-91
		CA-A- 2029372	08-05-91
		JP-A- 3188077	16-08-91
		US-A- 5100902	31-03-92